

表 1875 腎移植の回数 Body Mass Index別（腹膜透析患者）

Body Mass Index	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
14未満	22					22		5	27
(%)	(100.0)					(100.0)			
14～	66	1				67		4	71
(%)	(98.5)	(1.5)				(100.0)			
16～	203	2	1			206		22	228
(%)	(98.5)	(1.0)	(0.5)			(100.0)			
18～	554	13				567	1	59	627
(%)	(97.7)	(2.3)				(100.0)			
20～	1,027	10				1,037		103	1,140
(%)	(99.0)	(1.0)				(100.0)			
22～	1,103	7				1,110	4	147	1,261
(%)	(99.4)	(0.6)				(100.0)			
24～	814	3				817	2	89	908
(%)	(99.6)	(0.4)				(100.0)			
26～	496	2				498	2	61	561
(%)	(99.6)	(0.4)				(100.0)			
28～	267	3				270		32	302
(%)	(98.9)	(1.1)				(100.0)			
30～	192					192		28	220
(%)	(100.0)					(100.0)			
合計	4,744	41	1			4,786	9	550	5,345
(%)	(99.1)	(0.9)	(0.0)			(100.0)			
記載なし	1,031	15	1	1		1,048	14	2,534	3,596
(%)	(98.4)	(1.4)	(0.1)	(0.1)		(100.0)			
総計	5,775	56	2	1		5,834	23	3,084	8,941
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	23.14	21.37	16.98			23.12	23.64	23.33	23.15
標準偏差	3.59	3.17				3.59	2.22	3.68	3.60

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査