

表 1881 腎移植の回数 リン濃度(mg/dL)別（腹膜透析患者）

リン濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
3未満	101					101		25	126
(%)	(100.0)					(100.0)			
3.0～	236	3				239		31	270
(%)	(98.7)	(1.3)				(100.0)			
3.5～	457	4				461		64	525
(%)	(99.1)	(0.9)				(100.0)			
4.0～	670					670	2	98	770
(%)	(100.0)					(100.0)			
4.5～	810	4				814	1	106	921
(%)	(99.5)	(0.5)				(100.0)			
5.0～	831	8	1			840	1	107	948
(%)	(98.9)	(1.0)	(0.1)			(100.0)			
5.5～	715	16				731	1	107	839
(%)	(97.8)	(2.2)				(100.0)			
6.0～	119					119		19	138
(%)	(100.0)					(100.0)			
6.1～	405	3	1			409	3	65	477
(%)	(99.0)	(0.7)	(0.2)			(100.0)			
6.5～	338	3				341		58	399
(%)	(99.1)	(0.9)				(100.0)			
7.0～	183	1				184	1	27	212
(%)	(99.5)	(0.5)				(100.0)			
7.5～	127	2				129	1	19	149
(%)	(98.4)	(1.6)				(100.0)			
8.0～	63					63		8	71
(%)	(100.0)					(100.0)			
8.5～	30	2				32		6	38
(%)	(93.8)	(6.3)				(100.0)			
9.0～	76	1				77		12	89
(%)	(98.7)	(1.3)				(100.0)			
合計	5,161	47	2			5,210	10	752	5,972
(%)	(99.1)	(0.9)	(0.0)			(100.0)			
記載なし	614	9		1		624	13	2,332	2,969
(%)	(98.4)	(1.4)		(0.2)		(100.0)			
総計	5,775	56	2	1		5,834	23	3,084	8,941
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	5.24	5.61	5.70			5.24	5.77	5.26	5.25
標準偏差	1.34	1.40	0.85			1.34	1.23	1.41	1.35

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査