

表 1882 腎移植の回数 PTH値(pg/mL)別（腹膜透析患者）PTH測定法:intact-PTH

PTH値	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
20未満	105		1			106		17	123
(%)	(99.1)		(0.9)			(100.0)			
20～	250	2	1			253		28	281
(%)	(98.8)	(0.8)	(0.4)			(100.0)			
40～	236	4				240		28	268
(%)	(98.3)	(1.7)				(100.0)			
60～	277	2				279		36	315
(%)	(99.3)	(0.7)				(100.0)			
80～	283					283		26	309
(%)	(100.0)					(100.0)			
100～	280	3				283		33	316
(%)	(98.9)	(1.1)				(100.0)			
120～	265	2				267		38	305
(%)	(99.3)	(0.7)				(100.0)			
140～	247	3				250	3	32	285
(%)	(98.8)	(1.2)				(100.0)			
160～	247	1				248	1	27	276
(%)	(99.6)	(0.4)				(100.0)			
180～	572	11				583		71	654
(%)	(98.1)	(1.9)				(100.0)			
240～	8					8		2	10
(%)	(100.0)					(100.0)			
241～	382					382		49	431
(%)	(100.0)					(100.0)			
300～	290	4				294		37	331
(%)	(98.6)	(1.4)				(100.0)			
360～	170	1				171		29	200
(%)	(99.4)	(0.6)				(100.0)			
420～	118	2				120		19	139
(%)	(98.3)	(1.7)				(100.0)			
480～	38					38		5	43
(%)	(100.0)					(100.0)			
500～	45	1				46		8	54
(%)	(97.8)	(2.2)				(100.0)			
540～	60	3				63		10	73
(%)	(95.2)	(4.8)				(100.0)			
600～	92	1				93		12	105
(%)	(98.9)	(1.1)				(100.0)			
800～	89	2				91		19	110
(%)	(97.8)	(2.2)				(100.0)			
合計	4,054	42	2			4,098	4	526	4,628
(%)	(98.9)	(1.0)	(0.0)			(100.0)			
記載なし	298	2		1		301		75	376
(%)	(99.0)	(0.7)		(0.3)		(100.0)			
総計	4,352	44	2	1		4,399	4	601	5,004
(%)	(98.9)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	220.36	275.40	22.00			220.82	151.25	248.20	223.88
標準偏差	218.21	252.30	24.04			218.62	8.22	299.58	229.31

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査