

表 1883 腎移植の回数 PTH値(pg/mL)別（腹膜透析患者）PTH測定法:whole-PTH

| PTH値 | なし      | 1回     | 2回 | 3回 | 4回以上 | 合計      | 不明 | 記載なし   | 総計     |
|------|---------|--------|----|----|------|---------|----|--------|--------|
| 20未満 | 38      |        |    |    |      | 38      |    | 2      | 40     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 20～  | 40      |        |    |    |      | 40      |    | 5      | 45     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 35～  | 51      |        |    |    |      | 51      |    | 4      | 55     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 50～  | 66      |        |    |    |      | 66      |    | 11     | 77     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 75～  | 60      |        |    |    |      | 60      |    | 9      | 69     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 100～ | 63      |        |    |    |      | 63      |    | 8      | 71     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 125～ | 42      |        |    |    |      | 42      |    | 3      | 45     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 150～ | 1       |        |    |    |      | 1       |    |        | 1      |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 151～ | 36      |        |    |    |      | 36      |    | 7      | 43     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 180～ | 31      |        |    |    |      | 31      |    | 3      | 34     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 210～ | 22      | 1      |    |    |      | 23      |    | 4      | 27     |
| (%)  | (95.7)  | (4.3)  |    |    |      | (100.0) |    |        |        |
| 240～ | 21      | 1      |    |    |      | 22      |    | 1      | 23     |
| (%)  | (95.5)  | (4.5)  |    |    |      | (100.0) |    |        |        |
| 300～ | 19      |        |    |    |      | 19      |    |        | 19     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 350～ | 8       |        |    |    |      | 8       |    |        | 8      |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 400～ | 11      |        |    |    |      | 11      |    |        | 11     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 500～ | 10      |        |    |    |      | 10      |    | 1      | 11     |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 合計   | 519     | 2      |    |    |      | 521     |    | 58     | 579    |
| (%)  | (99.6)  | (0.4)  |    |    |      | (100.0) |    |        |        |
| 記載なし | 17      |        |    |    |      | 17      |    | 123    | 140    |
| (%)  | (100.0) |        |    |    |      | (100.0) |    |        |        |
| 総計   | 536     | 2      |    |    |      | 538     |    | 181    | 719    |
| (%)  | (99.6)  | (0.4)  |    |    |      | (100.0) |    |        |        |
| 平均   | 134.23  | 244.50 |    |    |      | 134.65  |    | 113.74 | 132.56 |
| 標準偏差 | 132.73  | 16.26  |    |    |      | 132.65  |    | 94.03  | 129.39 |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査