

表 1884 腎移植の回数 ヘモグロビン濃度(g/dL)別（腹膜透析患者）

ヘモグロビン 濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
5未満	6					6		1	7
(%)	(100.0)					(100.0)			
5～	13					13		1	14
(%)	(100.0)					(100.0)			
6～	24					24		4	28
(%)	(100.0)					(100.0)			
7～	100	3				103		21	124
(%)	(97.1)	(2.9)				(100.0)			
8～	279	7				286	1	57	344
(%)	(97.6)	(2.4)				(100.0)			
9～	792	5	1			798	3	102	903
(%)	(99.2)	(0.6)	(0.1)			(100.0)			
10～	1,480	15				1,495	4	207	1,706
(%)	(99.0)	(1.0)				(100.0)			
11～	1,498	7				1,505	2	202	1,709
(%)	(99.5)	(0.5)				(100.0)			
12～	674	9	1			684	1	111	796
(%)	(98.5)	(1.3)	(0.1)			(100.0)			
13～	203	2				205		32	237
(%)	(99.0)	(1.0)				(100.0)			
14～	69					69		10	79
(%)	(100.0)					(100.0)			
合計	5,138	48	2			5,188	11	748	5,947
(%)	(99.0)	(0.9)	(0.0)			(100.0)			
記載なし	637	8		1		646	12	2,336	2,994
(%)	(98.6)	(1.2)		(0.2)		(100.0)			
総計	5,775	56	2	1		5,834	23	3,084	8,941
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	10.82	10.53	11.05			10.82	10.35	10.79	10.81
標準偏差	1.41	1.62	2.33			1.41	1.03	1.45	1.42

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査