

表 1890 腎移植の回数 脈拍(拍/分)別 (腹膜透析患者)

脈拍	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
40未満	4					4		1	5
(%)	(100.0)					(100.0)			
40～	22					22		3	25
(%)	(100.0)					(100.0)			
50～	171	1				172		15	187
(%)	(99.4)	(0.6)				(100.0)			
60～	763	10				773	1	76	850
(%)	(98.7)	(1.3)				(100.0)			
70～	1,135	8				1,143	3	125	1,271
(%)	(99.3)	(0.7)				(100.0)			
80～	904	5				909		115	1,024
(%)	(99.4)	(0.6)				(100.0)			
90～	423	5				428	1	52	481
(%)	(98.8)	(1.2)				(100.0)			
100～	178	2	1			181		23	204
(%)	(98.3)	(1.1)	(0.6)			(100.0)			
110～	46					46		6	52
(%)	(100.0)					(100.0)			
120～	20					20			20
(%)	(100.0)					(100.0)			
合計	3,666	31	1			3,698	5	416	4,119
(%)	(99.1)	(0.8)	(0.0)			(100.0)			
記載なし	2,109	25	1	1		2,136	18	2,668	4,822
(%)	(98.7)	(1.2)	(0.0)	(0.0)		(100.0)			
総計	5,775	56	2	1		5,834	23	3,084	8,941
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	77.99	78.13	100.00			78.00	76.60	78.89	78.09
標準偏差	13.30	13.64				13.31	8.50	12.93	13.27

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査