

表 1894 透析時間(時間/回) 原疾患別(施設血液透析患者 透析回数週3回)

原疾患	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	148	5,576	4,479	46,766	4,492	5,817	291	371	23	178	68,141	436	68,577	4.04	0.53
(%)	(0.2)	(8.2)	(6.6)	(68.6)	(6.6)	(8.5)	(0.4)	(0.5)	(0.0)	(0.3)	(100.0)				
慢性腎盂腎炎	3	254	171	1,433	101	155	7	22		4	2,150	15	2,165	3.98	0.55
(%)	(0.1)	(11.8)	(8.0)	(66.7)	(4.7)	(7.2)	(0.3)	(1.0)		(0.2)	(100.0)				
急速進行性糸球体腎炎	5	223	159	1,242	68	114	4	9		2	1,826	8	1,834	3.94	0.51
(%)	(0.3)	(12.2)	(8.7)	(68.0)	(3.7)	(6.2)	(0.2)	(0.5)		(0.1)	(100.0)				
妊娠腎/妊娠中毒症	5	58	85	746	85	73	5	2		4	1,063	1	1,064	4.04	0.50
(%)	(0.5)	(5.5)	(8.0)	(70.2)	(8.0)	(6.9)	(0.5)	(0.2)		(0.4)	(100.0)				
その他の分類不能の腎炎	2	95	64	624	64	82	3	2	1	9	946	3	949	4.04	0.63
(%)	(0.2)	(10.0)	(6.8)	(66.0)	(6.8)	(8.7)	(0.3)	(0.2)	(0.1)	(1.0)	(100.0)				
多発性嚢胞腎	12	793	670	5,595	456	547	14	28		29	8,144	54	8,198	3.99	0.52
(%)	(0.1)	(9.7)	(8.2)	(68.7)	(5.6)	(6.7)	(0.2)	(0.3)		(0.4)	(100.0)				
腎硬化症	53	3,202	2,098	13,930	745	1,160	25	89	7	51	21,360	100	21,460	3.90	0.54
(%)	(0.2)	(15.0)	(9.8)	(65.2)	(3.5)	(5.4)	(0.1)	(0.4)	(0.0)	(0.2)	(100.0)				
悪性高血圧	3	217	152	1,226	115	147	5	7		3	1,875	6	1,881	3.98	0.52
(%)	(0.2)	(11.6)	(8.1)	(65.4)	(6.1)	(7.8)	(0.3)	(0.4)		(0.2)	(100.0)				
糖尿病性腎症	178	10,086	7,241	61,868	4,114	5,838	168	362	18	156	90,029	478	90,507	3.96	0.51
(%)	(0.2)	(11.2)	(8.0)	(68.7)	(4.6)	(6.5)	(0.2)	(0.4)	(0.0)	(0.2)	(100.0)				
SLE腎炎	3	170	132	1,042	80	100	3	7		4	1,541	9	1,550	3.97	0.53
(%)	(0.2)	(11.0)	(8.6)	(67.6)	(5.2)	(6.5)	(0.2)	(0.5)		(0.3)	(100.0)				
アミロイド腎	4	39	38	199	20	23	1				324	1	325	3.92	0.52
(%)	(1.2)	(12.0)	(11.7)	(61.4)	(6.2)	(7.1)	(0.3)				(100.0)				
痛風腎	1	73	65	574	42	63	1	3		4	826	2	828	4.01	0.52
(%)	(0.1)	(8.8)	(7.9)	(69.5)	(5.1)	(7.6)	(0.1)	(0.4)		(0.5)	(100.0)				
先天性代謝異常による腎不全	1	14	14	118	10	17	1				175	2	177	4.01	0.50
(%)	(0.6)	(8.0)	(8.0)	(67.4)	(5.7)	(9.7)	(0.6)				(100.0)				
腎・尿路結核	1	19	13	106	7	11	1				158		158	3.94	0.49
(%)	(0.6)	(12.0)	(8.2)	(67.1)	(4.4)	(7.0)	(0.6)				(100.0)				
腎・尿路結石	1	54	36	295	18	21	2	2		1	430		430	3.93	0.51
(%)	(0.2)	(12.6)	(8.4)	(68.6)	(4.2)	(4.9)	(0.5)	(0.5)		(0.2)	(100.0)				
腎・尿路腫瘍	1	79	58	485	19	33	1	4		2	682	4	686	3.94	0.51
(%)	(0.1)	(11.6)	(8.5)	(71.1)	(2.8)	(4.8)	(0.1)	(0.6)		(0.3)	(100.0)				
閉塞性尿路障害	2	70	40	332	25	37	1			5	512	5	517	3.97	0.64
(%)	(0.4)	(13.7)	(7.8)	(64.8)	(4.9)	(7.2)	(0.2)			(1.0)	(100.0)				
骨髄腫	2	51	20	99	5	8	1		1	1	188	2	190	3.76	0.68
(%)	(1.1)	(27.1)	(10.6)	(52.7)	(2.7)	(4.3)	(0.5)		(0.5)	(0.5)	(100.0)				
腎形成不全		31	24	281	37	37	1	2		2	415	4	419	4.07	0.55
(%)		(7.5)	(5.8)	(67.7)	(8.9)	(8.9)	(0.2)	(0.5)		(0.5)	(100.0)				
不明	77	2,876	1,829	13,448	819	870	34	46	4	27	20,030	95	20,125	3.89	0.50
(%)	(0.4)	(14.4)	(9.1)	(67.1)	(4.1)	(4.3)	(0.2)	(0.2)	(0.0)	(0.1)	(100.0)				
再導入	1	111	82	944	104	139	5	9	3	7	1,405	12	1,417	4.08	0.58
(%)	(0.1)	(7.9)	(5.8)	(67.2)	(7.4)	(9.9)	(0.4)	(0.6)	(0.2)	(0.5)	(100.0)				
その他	18	729	469	3,645	252	295	21	29	1	22	5,481	33	5,514	3.94	0.56
(%)	(0.3)	(13.3)	(8.6)	(66.5)	(4.6)	(5.4)	(0.4)	(0.5)	(0.0)	(0.4)	(100.0)				
合計	521	24,820	17,939	154,998	11,678	15,587	595	994	58	511	227,701	1,270	228,971	3.97	0.52
(%)	(0.2)	(10.9)	(7.9)	(68.1)	(5.1)	(6.8)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
記載なし															
(%)															
総計	521	24,820	17,939	154,998	11,678	15,587	595	994	58	511	227,701	1,270	228,971	3.97	0.52
(%)	(0.2)	(10.9)	(7.9)	(68.1)	(5.1)	(6.8)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査