

表 1899 透析時間(時間/回) 体重減少量(kg)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

体重減少量	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
0.0未満	3	197	61	454	14	23		3			755		755	3.75	0.53
(%)	(0.4)	(26.1)	(8.1)	(60.1)	(1.9)	(3.0)		(0.4)			(100.0)				
0.0～	67	2,015	857	4,931	172	320	16	35	2	13	8,428	22	8,450	3.77	0.56
(%)	(0.8)	(23.9)	(10.2)	(58.5)	(2.0)	(3.8)	(0.2)	(0.4)	(0.0)	(0.2)	(100.0)				
1.0～	133	5,429	4,305	27,351	1,378	2,028	88	177	5	38	40,932	93	41,025	3.90	0.50
(%)	(0.3)	(13.3)	(10.5)	(66.8)	(3.4)	(5.0)	(0.2)	(0.4)	(0.0)	(0.1)	(100.0)				
2.0～	66	4,216	5,868	53,486	3,846	4,889	193	295	21	127	73,007	154	73,161	4.02	0.46
(%)	(0.1)	(5.8)	(8.0)	(73.3)	(5.3)	(6.7)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
3.0～	15	673	1,569	28,283	3,251	3,774	144	194	12	132	38,047	63	38,110	4.14	0.46
(%)	(0.0)	(1.8)	(4.1)	(74.3)	(8.5)	(9.9)	(0.4)	(0.5)	(0.0)	(0.3)	(100.0)				
4.0～	4	77	189	5,835	1,295	1,769	60	95	11	98	9,433	12	9,445	4.32	0.60
(%)	(0.0)	(0.8)	(2.0)	(61.9)	(13.7)	(18.8)	(0.6)	(1.0)	(0.1)	(1.0)	(100.0)				
5.0～		17	32	926	261	606	36	52	1	22	1,953	5	1,958	4.50	0.67
(%)		(0.9)	(1.6)	(47.4)	(13.4)	(31.0)	(1.8)	(2.7)	(0.1)	(1.1)	(100.0)				
6.0～		1	5	168	57	130	12	12	2	8	395		395	4.60	0.73
(%)		(0.3)	(1.3)	(42.5)	(14.4)	(32.9)	(3.0)	(3.0)	(0.5)	(2.0)	(100.0)				
7.0～			3	36	12	32	3	5		4	95		95	4.70	0.85
(%)			(3.2)	(37.9)	(12.6)	(33.7)	(3.2)	(5.3)		(4.2)	(100.0)				
8.0～	1	10	7	108	8	20	1	3		2	160		160	4.16	0.77
(%)	(0.6)	(6.3)	(4.4)	(67.5)	(5.0)	(12.5)	(0.6)	(1.9)		(1.3)	(100.0)				
合計	289	12,635	12,896	121,578	10,294	13,591	553	871	54	444	173,205	349	173,554	4.03	0.51
(%)	(0.2)	(7.3)	(7.4)	(70.2)	(5.9)	(7.8)	(0.3)	(0.5)	(0.0)	(0.3)	(100.0)				
記載なし	6	530	271	2,806	161	195	3	4		1	3,977	678	4,655	3.91	0.47
(%)	(0.2)	(13.3)	(6.8)	(70.6)	(4.0)	(4.9)	(0.1)	(0.1)		(0.0)	(100.0)				
総計	295	13,165	13,167	124,384	10,455	13,786	556	875	54	445	177,182	1,027	178,209	4.03	0.50
(%)	(0.2)	(7.4)	(7.4)	(70.2)	(5.9)	(7.8)	(0.3)	(0.5)	(0.0)	(0.3)	(100.0)				
平均	1.62	1.72	2.13	2.50	2.95	2.98	3.02	2.88	3.08	3.35	2.49	2.37	2.49		
標準偏差	1.29	0.93	0.88	0.99	1.07	1.20	1.29	1.35	1.19	1.58	1.05	0.96	1.05		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査