

表 1900 透析時間(時間/回) 体重減少率(%)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

体重減少率	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
1.0未満	36	971	351	2,083	70	145	5	17		6	3,684	12	3,696	3.75	0.58
(%)	(1.0)	(26.4)	(9.5)	(56.5)	(1.9)	(3.9)	(0.1)	(0.5)		(0.2)	(100.0)				
1.0～	33	1,216	603	4,162	191	322	17	37	3	10	6,594	15	6,609	3.86	0.55
(%)	(0.5)	(18.4)	(9.1)	(63.1)	(2.9)	(4.9)	(0.3)	(0.6)	(0.0)	(0.2)	(100.0)				
2.0～	56	2,240	1,676	11,145	652	947	48	92	2	21	16,879	42	16,921	3.92	0.52
(%)	(0.3)	(13.3)	(9.9)	(66.0)	(3.9)	(5.6)	(0.3)	(0.5)	(0.0)	(0.1)	(100.0)				
3.0～	64	3,118	3,061	23,896	1,667	2,390	95	157	9	53	34,510	74	34,584	3.98	0.49
(%)	(0.2)	(9.0)	(8.9)	(69.2)	(4.8)	(6.9)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
4.0～	56	2,726	3,436	33,306	2,815	3,545	130	200	15	106	46,335	80	46,415	4.04	0.48
(%)	(0.1)	(5.9)	(7.4)	(71.9)	(6.1)	(7.7)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
5.0～	24	1,471	2,297	26,372	2,554	3,078	130	168	12	103	36,209	80	36,289	4.08	0.47
(%)	(0.1)	(4.1)	(6.3)	(72.8)	(7.1)	(8.5)	(0.4)	(0.5)	(0.0)	(0.3)	(100.0)				
6.0～	14	615	1,015	13,285	1,419	1,790	63	109	9	75	18,394	33	18,427	4.12	0.50
(%)	(0.1)	(3.3)	(5.5)	(72.2)	(7.7)	(9.7)	(0.3)	(0.6)	(0.0)	(0.4)	(100.0)				
7.0～	2	193	312	4,998	641	824	37	48	3	44	7,102	6	7,108	4.17	0.54
(%)	(0.0)	(2.7)	(4.4)	(70.4)	(9.0)	(11.6)	(0.5)	(0.7)	(0.0)	(0.6)	(100.0)				
8.0～	1	52	105	1,478	175	320	18	28		13	2,190	5	2,195	4.21	0.57
(%)	(0.0)	(2.4)	(4.8)	(67.5)	(8.0)	(14.6)	(0.8)	(1.3)		(0.6)	(100.0)				
9.0～	2	23	34	779	105	222	10	15	1	12	1,203	2	1,205	4.28	0.62
(%)	(0.2)	(1.9)	(2.8)	(64.8)	(8.7)	(18.5)	(0.8)	(1.2)	(0.1)	(1.0)	(100.0)				
合計	288	12,625	12,890	121,504	10,289	13,583	553	871	54	443	173,100	349	173,449	4.03	0.51
(%)	(0.2)	(7.3)	(7.4)	(70.2)	(5.9)	(7.8)	(0.3)	(0.5)	(0.0)	(0.3)	(100.0)				
記載なし	7	540	277	2,880	166	203	3	4		2	4,082	678	4,760	3.91	0.47
(%)	(0.2)	(13.2)	(6.8)	(70.6)	(4.1)	(5.0)	(0.1)	(0.1)		(0.0)	(100.0)				
総計	295	13,165	13,167	124,384	10,455	13,786	556	875	54	445	177,182	1,027	178,209	4.03	0.50
(%)	(0.2)	(7.4)	(7.4)	(70.2)	(5.9)	(7.8)	(0.3)	(0.5)	(0.0)	(0.3)	(100.0)				
平均	3.28	3.56	4.20	4.59	4.96	4.93	4.94	4.74	4.93	5.28	4.53	4.33	4.53		
標準偏差	1.81	1.72	1.58	1.60	1.55	1.69	1.80	1.88	1.58	1.77	1.65	1.65	1.65		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査