

表 1911 透析時間(時間/回) 透析前リン濃度(mg/dL)別(施設血液透析患者 透析回数週3回)

透析前リン濃度	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
3未満	50	1,536	691	5,728	311	526	17	41	1	25	8,926	35	8,961	3.89	0.57
(%)	(0.6)	(17.2)	(7.7)	(64.2)	(3.5)	(5.9)	(0.2)	(0.5)	(0.0)	(0.3)	(100.0)				
3.0～	47	1,531	853	7,268	417	740	22	46	2	31	10,957	23	10,980	3.93	0.56
(%)	(0.4)	(14.0)	(7.8)	(66.3)	(3.8)	(6.8)	(0.2)	(0.4)	(0.0)	(0.3)	(100.0)				
3.5～	51	2,363	1,468	12,714	870	1,287	50	113	6	56	18,978	40	19,018	3.96	0.55
(%)	(0.3)	(12.5)	(7.7)	(67.0)	(4.6)	(6.8)	(0.3)	(0.6)	(0.0)	(0.3)	(100.0)				
4.0～	70	3,240	2,285	18,933	1,350	1,943	79	141	7	68	28,116	76	28,192	3.97	0.54
(%)	(0.2)	(11.5)	(8.1)	(67.3)	(4.8)	(6.9)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
4.5～	52	3,564	2,601	22,730	1,732	2,340	93	173	5	78	33,368	100	33,468	3.98	0.52
(%)	(0.2)	(10.7)	(7.8)	(68.1)	(5.2)	(7.0)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
5.0～	62	3,317	2,760	23,175	1,751	2,336	98	143	11	71	33,724	85	33,809	3.98	0.51
(%)	(0.2)	(9.8)	(8.2)	(68.7)	(5.2)	(6.9)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
5.5～	56	2,701	2,269	19,992	1,619	2,063	76	103	7	53	28,939	73	29,012	3.99	0.50
(%)	(0.2)	(9.3)	(7.8)	(69.1)	(5.6)	(7.1)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
6.0～	5	442	387	3,414	284	337	15	12	2	11	4,909	19	4,928	3.99	0.49
(%)	(0.1)	(9.0)	(7.9)	(69.5)	(5.8)	(6.9)	(0.3)	(0.2)	(0.0)	(0.2)	(100.0)				
6.1～	28	1,553	1,314	11,327	906	1,158	48	55	5	39	16,433	72	16,505	3.99	0.51
(%)	(0.2)	(9.5)	(8.0)	(68.9)	(5.5)	(7.0)	(0.3)	(0.3)	(0.0)	(0.2)	(100.0)				
6.5～	31	1,340	1,120	9,901	814	973	41	54	1	39	14,314	63	14,377	3.99	0.51
(%)	(0.2)	(9.4)	(7.8)	(69.2)	(5.7)	(6.8)	(0.3)	(0.4)	(0.0)	(0.3)	(100.0)				
7.0～	17	756	677	6,184	527	585	19	44	5	17	8,831	29	8,860	4.00	0.50
(%)	(0.2)	(8.6)	(7.7)	(70.0)	(6.0)	(6.6)	(0.2)	(0.5)	(0.1)	(0.2)	(100.0)				
7.5～	12	477	376	3,720	339	369	17	29	1	5	5,345	14	5,359	4.00	0.49
(%)	(0.2)	(8.9)	(7.0)	(69.6)	(6.3)	(6.9)	(0.3)	(0.5)	(0.0)	(0.1)	(100.0)				
8.0～	10	314	226	2,110	221	268	9	10	3	7	3,178	14	3,192	4.01	0.53
(%)	(0.3)	(9.9)	(7.1)	(66.4)	(7.0)	(8.4)	(0.3)	(0.3)	(0.1)	(0.2)	(100.0)				
8.5～	2	190	153	1,394	124	142	1	6	1	4	2,017	6	2,023	3.99	0.49
(%)	(0.1)	(9.4)	(7.6)	(69.1)	(6.1)	(7.0)	(0.0)	(0.3)	(0.0)	(0.2)	(100.0)				
9.0～	8	312	207	2,090	202	300	5	14	1	5	3,144	12	3,156	4.02	0.53
(%)	(0.3)	(9.9)	(6.6)	(66.5)	(6.4)	(9.5)	(0.2)	(0.4)	(0.0)	(0.2)	(100.0)				
合計	501	23,636	17,387	150,680	11,467	15,367	590	984	58	509	221,179	661	221,840	3.98	0.52
(%)	(0.2)	(10.7)	(7.9)	(68.1)	(5.2)	(6.9)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
記載なし	20	1,184	552	4,318	211	220	5	10		2	6,522	609	7,131	3.84	0.49
(%)	(0.3)	(18.2)	(8.5)	(66.2)	(3.2)	(3.4)	(0.1)	(0.2)		(0.0)	(100.0)				
総計	521	24,820	17,939	154,998	11,678	15,587	595	994	58	511	227,701	1,270	228,971	3.97	0.52
(%)	(0.2)	(10.9)	(7.9)	(68.1)	(5.1)	(6.8)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
平均	4.86	4.99	5.17	5.22	5.37	5.26	5.21	5.08	5.54	5.04	5.20	5.35	5.20		
標準偏差	1.62	1.49	1.40	1.41	1.42	1.46	1.33	1.42	1.71	1.36	1.43	1.51	1.43		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査