

表 1914 透析時間(時間/回) 透析前ヘモグロビン濃度(g/dL)別 (施設血液透析患者 透析回数週3回)

透析前ヘモグロビン濃度	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
5未満	1	28	18	200	7	6					260	1	261	3.89	0.40
(%)	(0.4)	(10.8)	(6.9)	(76.9)	(2.7)	(2.3)					(100.0)				
5～	2	69	40	233	8	7		1			360	2	362	3.80	0.47
(%)	(0.6)	(19.2)	(11.1)	(64.7)	(2.2)	(1.9)		(0.3)			(100.0)				
6～	8	214	73	567	30	37	2			1	932	10	942	3.79	0.54
(%)	(0.9)	(23.0)	(7.8)	(60.8)	(3.2)	(4.0)	(0.2)			(0.1)	(100.0)				
7～	26	738	276	2,262	100	142	5	7		11	3,567	11	3,578	3.82	0.56
(%)	(0.7)	(20.7)	(7.7)	(63.4)	(2.8)	(4.0)	(0.1)	(0.2)		(0.3)	(100.0)				
8～	63	2,022	929	7,660	407	533	19	23	3	30	11,689	37	11,726	3.86	0.54
(%)	(0.5)	(17.3)	(7.9)	(65.5)	(3.5)	(4.6)	(0.2)	(0.2)	(0.0)	(0.3)	(100.0)				
9～	97	4,785	3,110	25,077	1,532	2,001	87	149	11	102	36,951	87	37,038	3.93	0.53
(%)	(0.3)	(12.9)	(8.4)	(67.9)	(4.1)	(5.4)	(0.2)	(0.4)	(0.0)	(0.3)	(100.0)				
10～	159	7,463	6,103	51,828	3,975	4,861	189	293	18	131	75,020	197	75,217	3.98	0.50
(%)	(0.2)	(9.9)	(8.1)	(69.1)	(5.3)	(6.5)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
11～	89	5,659	4,763	41,963	3,408	4,833	169	295	13	147	61,339	140	61,479	4.01	0.52
(%)	(0.1)	(9.2)	(7.8)	(68.4)	(5.6)	(7.9)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
12～	41	1,979	1,595	15,005	1,396	2,016	80	135	10	57	22,314	56	22,370	4.03	0.54
(%)	(0.2)	(8.9)	(7.1)	(67.2)	(6.3)	(9.0)	(0.4)	(0.6)	(0.0)	(0.3)	(100.0)				
13～	14	494	317	3,694	359	548	26	47	3	20	5,522	15	5,537	4.06	0.57
(%)	(0.3)	(8.9)	(5.7)	(66.9)	(6.5)	(9.9)	(0.5)	(0.9)	(0.1)	(0.4)	(100.0)				
14～	7	123	100	1,253	124	257	11	35		10	1,920	13	1,933	4.14	0.62
(%)	(0.4)	(6.4)	(5.2)	(65.3)	(6.5)	(13.4)	(0.6)	(1.8)		(0.5)	(100.0)				
合計	507	23,574	17,324	149,742	11,346	15,241	588	985	58	509	219,874	569	220,443	3.98	0.52
(%)	(0.2)	(10.7)	(7.9)	(68.1)	(5.2)	(6.9)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
記載なし	14	1,246	615	5,256	332	346	7	9		2	7,827	701	8,528	3.88	0.48
(%)	(0.2)	(15.9)	(7.9)	(67.2)	(4.2)	(4.4)	(0.1)	(0.1)		(0.0)	(100.0)				
総計	521	24,820	17,939	154,998	11,678	15,587	595	994	58	511	227,701	1,270	228,971	3.97	0.52
(%)	(0.2)	(10.9)	(7.9)	(68.1)	(5.1)	(6.8)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
平均	10.25	10.41	10.62	10.69	10.88	10.97	10.96	11.10	10.91	10.77	10.69	10.62	10.69		
標準偏差	1.57	1.37	1.23	1.26	1.21	1.26	1.28	1.38	1.31	1.40	1.28	1.47	1.28		

数値下のカッコ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査