

表 1915 透析時間(時間/回) 総コレステロール濃度(mg/dL)別 (施設血液透析患者 透析回数週3回)

総コレステロール濃度	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
100未満	18	785	509	4,392	288	390	13	38		7	6,440	14	6,454	3.95	0.51
(%)	(0.3)	(12.2)	(7.9)	(68.2)	(4.5)	(6.1)	(0.2)	(0.6)		(0.1)	(100.0)				
100～	40	1,968	1,469	12,742	965	1,258	40	74	3	26	18,585	35	18,620	3.97	0.50
(%)	(0.2)	(10.6)	(7.9)	(68.6)	(5.2)	(6.8)	(0.2)	(0.4)	(0.0)	(0.1)	(100.0)				
120～	61	3,478	2,622	23,518	1,757	2,493	87	151	8	64	34,239	76	34,315	3.99	0.51
(%)	(0.2)	(10.2)	(7.7)	(68.7)	(5.1)	(7.3)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
140～	75	3,873	3,005	26,508	2,053	2,819	101	204	13	86	38,737	89	38,826	3.99	0.52
(%)	(0.2)	(10.0)	(7.8)	(68.4)	(5.3)	(7.3)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
160～	69	3,097	2,474	21,337	1,722	2,471	88	172	11	81	31,522	61	31,583	4.00	0.53
(%)	(0.2)	(9.8)	(7.8)	(67.7)	(5.5)	(7.8)	(0.3)	(0.5)	(0.0)	(0.3)	(100.0)				
180～	53	1,942	1,524	13,199	1,074	1,545	61	110	7	61	19,576	43	19,619	4.00	0.54
(%)	(0.3)	(9.9)	(7.8)	(67.4)	(5.5)	(7.9)	(0.3)	(0.6)	(0.0)	(0.3)	(100.0)				
200～	27	1,035	788	6,864	548	790	40	59	2	31	10,184	25	10,209	4.00	0.55
(%)	(0.3)	(10.2)	(7.7)	(67.4)	(5.4)	(7.8)	(0.4)	(0.6)	(0.0)	(0.3)	(100.0)				
220～	15	538	365	2,912	238	335	15	25	3	23	4,469	6	4,475	3.98	0.58
(%)	(0.3)	(12.0)	(8.2)	(65.2)	(5.3)	(7.5)	(0.3)	(0.6)	(0.1)	(0.5)	(100.0)				
240～	4	239	159	1,085	80	144	3	7	1	8	1,730	3	1,733	3.96	0.59
(%)	(0.2)	(13.8)	(9.2)	(62.7)	(4.6)	(8.3)	(0.2)	(0.4)	(0.1)	(0.5)	(100.0)				
260～	6	198	82	689	36	86	2	7		3	1,109	2	1,111	3.90	0.60
(%)	(0.5)	(17.9)	(7.4)	(62.1)	(3.2)	(7.8)	(0.2)	(0.6)		(0.3)	(100.0)				
合計	368	17,153	12,997	113,246	8,761	12,331	450	847	48	390	166,591	354	166,945	3.99	0.53
(%)	(0.2)	(10.3)	(7.8)	(68.0)	(5.3)	(7.4)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
記載なし	153	7,667	4,942	41,752	2,917	3,256	145	147	10	121	61,110	916	62,026	3.93	0.51
(%)	(0.3)	(12.5)	(8.1)	(68.3)	(4.8)	(5.3)	(0.2)	(0.2)	(0.0)	(0.2)	(100.0)				
総計	521	24,820	17,939	154,998	11,678	15,587	595	994	58	511	227,701	1,270	228,971	3.97	0.52
(%)	(0.2)	(10.9)	(7.9)	(68.1)	(5.1)	(6.8)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
平均	159.10	155.20	154.93	154.28	155.34	156.56	158.04	157.43	163.63	165.55	154.72	154.39	154.71		
標準偏差	40.38	37.89	35.41	34.93	34.38	34.84	34.79	36.31	32.68	36.84	35.29	33.75	35.28		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査