

表 1916 透析時間(時間/回) HDL-C濃度(mg/dL)別 (施設血液透析患者 透析回数週3回)

HDL-C濃度	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
20未満	2	124	38	429	27	39	3	3		1	666	5	671	3.89	0.56
(%)	(0.3)	(18.6)	(5.7)	(64.4)	(4.1)	(5.9)	(0.5)	(0.5)		(0.2)	(100.0)				
20～	28	1,314	997	8,593	737	1,007	41	57	6	35	12,815	31	12,846	4.00	0.54
(%)	(0.2)	(10.3)	(7.8)	(67.1)	(5.8)	(7.9)	(0.3)	(0.4)	(0.0)	(0.3)	(100.0)				
30～	61	3,680	2,715	25,025	2,063	2,877	108	134	11	120	36,794	94	36,888	4.00	0.53
(%)	(0.2)	(10.0)	(7.4)	(68.0)	(5.6)	(7.8)	(0.3)	(0.4)	(0.0)	(0.3)	(100.0)				
40～	76	3,993	3,170	27,866	2,168	2,963	94	171	17	116	40,634	95	40,729	3.99	0.52
(%)	(0.2)	(9.8)	(7.8)	(68.6)	(5.3)	(7.3)	(0.2)	(0.4)	(0.0)	(0.3)	(100.0)				
50～	44	2,815	2,281	19,987	1,483	2,137	74	126	10	78	29,035	45	29,080	3.99	0.52
(%)	(0.2)	(9.7)	(7.9)	(68.8)	(5.1)	(7.4)	(0.3)	(0.4)	(0.0)	(0.3)	(100.0)				
60～	39	1,608	1,335	11,519	868	1,206	46	93	5	46	16,765	27	16,792	3.99	0.53
(%)	(0.2)	(9.6)	(8.0)	(68.7)	(5.2)	(7.2)	(0.3)	(0.6)	(0.0)	(0.3)	(100.0)				
70～	18	767	593	5,653	443	611	30	44		30	8,189	12	8,201	4.01	0.54
(%)	(0.2)	(9.4)	(7.2)	(69.0)	(5.4)	(7.5)	(0.4)	(0.5)		(0.4)	(100.0)				
80～	7	351	304	2,613	202	296	15	23	2	16	3,829	5	3,834	4.02	0.55
(%)	(0.2)	(9.2)	(7.9)	(68.2)	(5.3)	(7.7)	(0.4)	(0.6)	(0.1)	(0.4)	(100.0)				
90～	1	165	130	1,187	74	154	3	7		11	1,732	1	1,733	4.03	0.58
(%)	(0.1)	(9.5)	(7.5)	(68.5)	(4.3)	(8.9)	(0.2)	(0.4)		(0.6)	(100.0)				
100～	2	142	109	1,034	58	100	1	7		4	1,457	1	1,458	3.99	0.51
(%)	(0.1)	(9.7)	(7.5)	(71.0)	(4.0)	(6.9)	(0.1)	(0.5)		(0.3)	(100.0)				
合計	278	14,959	11,672	103,906	8,123	11,390	415	665	51	457	151,916	316	152,232	4.00	0.53
(%)	(0.2)	(9.8)	(7.7)	(68.4)	(5.3)	(7.5)	(0.3)	(0.4)	(0.0)	(0.3)	(100.0)				
記載なし	243	9,861	6,267	51,092	3,555	4,197	180	329	7	54	75,785	954	76,739	3.92	0.51
(%)	(0.3)	(13.0)	(8.3)	(67.4)	(4.7)	(5.5)	(0.2)	(0.4)	(0.0)	(0.1)	(100.0)				
総計	521	24,820	17,939	154,998	11,678	15,587	595	994	58	511	227,701	1,270	228,971	3.97	0.52
(%)	(0.2)	(10.9)	(7.9)	(68.1)	(5.1)	(6.8)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
平均	48.03	47.76	48.40	48.39	47.59	48.03	47.72	49.74	45.45	48.92	48.26	44.45	48.25		
標準偏差	17.07	16.51	16.28	16.61	16.03	16.55	16.32	17.48	14.02	17.14	16.55	14.60	16.54		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査