

表 1918 透析時間(時間/回) 透析前収縮期血圧(mmHg)別 (施設血液透析患者 透析回数週3回)

透析前収縮期血圧	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
60未満		7	2	22		3					34	1	35	3.86	0.55
(%)		(20.6)	(5.9)	(64.7)		(8.8)					(100.0)				
60～		29	25	175	15	19	2	5	1		271		271	4.01	0.57
(%)		(10.7)	(9.2)	(64.6)	(5.5)	(7.0)	(0.7)	(1.8)	(0.4)		(100.0)				
80～	15	301	190	1,541	112	182	10	19		6	2,376	1	2,377	3.96	0.57
(%)	(0.6)	(12.7)	(8.0)	(64.9)	(4.7)	(7.7)	(0.4)	(0.8)		(0.3)	(100.0)				
100～	57	1,681	1,025	8,767	601	876	36	69	1	30	13,143	12	13,155	3.95	0.54
(%)	(0.4)	(12.8)	(7.8)	(66.7)	(4.6)	(6.7)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
120～	96	4,827	3,200	28,033	2,087	2,835	123	204	17	102	41,524	52	41,576	3.97	0.53
(%)	(0.2)	(11.6)	(7.7)	(67.5)	(5.0)	(6.8)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
140～	137	6,686	5,059	43,722	3,347	4,567	133	280	18	163	64,112	91	64,203	3.98	0.52
(%)	(0.2)	(10.4)	(7.9)	(68.2)	(5.2)	(7.1)	(0.2)	(0.4)	(0.0)	(0.3)	(100.0)				
160～	110	4,600	3,724	32,101	2,482	3,287	124	222	9	108	46,767	76	46,843	3.99	0.52
(%)	(0.2)	(9.8)	(8.0)	(68.6)	(5.3)	(7.0)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
180～	42	1,755	1,423	12,203	994	1,364	45	82	6	49	17,963	39	18,002	4.00	0.53
(%)	(0.2)	(9.8)	(7.9)	(67.9)	(5.5)	(7.6)	(0.3)	(0.5)	(0.0)	(0.3)	(100.0)				
200～	7	400	329	3,081	262	366	19	30	3	7	4,504	11	4,515	4.02	0.52
(%)	(0.2)	(8.9)	(7.3)	(68.4)	(5.8)	(8.1)	(0.4)	(0.7)	(0.1)	(0.2)	(100.0)				
220～	1	85	66	586	51	85	4	9		5	892	2	894	4.05	0.59
(%)	(0.1)	(9.5)	(7.4)	(65.7)	(5.7)	(9.5)	(0.4)	(1.0)		(0.6)	(100.0)				
合計	465	20,371	15,043	130,231	9,951	13,584	496	920	55	470	191,586	285	191,871	3.98	0.53
(%)	(0.2)	(10.6)	(7.9)	(68.0)	(5.2)	(7.1)	(0.3)	(0.5)	(0.0)	(0.2)	(100.0)				
記載なし	56	4,449	2,896	24,767	1,727	2,003	99	74	3	41	36,115	985	37,100	3.93	0.49
(%)	(0.2)	(12.3)	(8.0)	(68.6)	(4.8)	(5.5)	(0.3)	(0.2)	(0.0)	(0.1)	(100.0)				
総計	521	24,820	17,939	154,998	11,678	15,587	595	994	58	511	227,701	1,270	228,971	3.97	0.52
(%)	(0.2)	(10.9)	(7.9)	(68.1)	(5.1)	(6.8)	(0.3)	(0.4)	(0.0)	(0.2)	(100.0)				
平均	147.71	149.59	151.76	151.78	152.77	152.47	151.58	151.22	153.07	152.12	151.63	156.88	151.64		
標準偏差	25.47	24.16	23.96	23.90	24.05	24.39	26.64	26.40	25.48	23.63	24.01	24.72	24.02		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査