

表 1923 透析時間(時間/回) 透析歴(年)別 (血液透析濾過患者 透析回数週3回)

透析歴	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
2年未満	29	836	480	3,125	221	208	7	19	1	2	4,928	36	4,964	3.87	0.52
(%)	(0.6)	(17.0)	(9.7)	(63.4)	(4.5)	(4.2)	(0.1)	(0.4)	(0.0)	(0.0)	(100.0)				
2年～	27	715	622	5,879	594	640	32	48	1	9	8,567	48	8,615	4.02	0.49
(%)	(0.3)	(8.3)	(7.3)	(68.6)	(6.9)	(7.5)	(0.4)	(0.6)	(0.0)	(0.1)	(100.0)				
5年～	16	444	763	6,832	944	957	46	48	4	6	10,060	56	10,116	4.09	0.47
(%)	(0.2)	(4.4)	(7.6)	(67.9)	(9.4)	(9.5)	(0.5)	(0.5)	(0.0)	(0.1)	(100.0)				
10年～	14	159	296	4,155	692	737	40	61	2	6	6,162	32	6,194	4.17	0.49
(%)	(0.2)	(2.6)	(4.8)	(67.4)	(11.2)	(12.0)	(0.6)	(1.0)	(0.0)	(0.1)	(100.0)				
15年～	3	56	127	2,603	507	525	38	32		7	3,898	26	3,924	4.22	0.48
(%)	(0.1)	(1.4)	(3.3)	(66.8)	(13.0)	(13.5)	(1.0)	(0.8)		(0.2)	(100.0)				
20年～		29	79	1,726	366	364	20	33	5	3	2,625	12	2,637	4.24	0.48
(%)		(1.1)	(3.0)	(65.8)	(13.9)	(13.9)	(0.8)	(1.3)	(0.2)	(0.1)	(100.0)				
25年～	2	22	46	1,018	234	280	23	17	2	2	1,646	5	1,651	4.28	0.50
(%)	(0.1)	(1.3)	(2.8)	(61.8)	(14.2)	(17.0)	(1.4)	(1.0)	(0.1)	(0.1)	(100.0)				
30年～		12	33	675	173	196	8	17	2		1,116	7	1,123	4.29	0.49
(%)		(1.1)	(3.0)	(60.5)	(15.5)	(17.6)	(0.7)	(1.5)	(0.2)		(100.0)				
35年～		7	20	353	93	101	2	9			585	2	587	4.27	0.48
(%)		(1.2)	(3.4)	(60.3)	(15.9)	(17.3)	(0.3)	(1.5)			(100.0)				
40年～	1	1	11	62	23	23	2			1	124		124	4.29	0.59
(%)	(0.8)	(0.8)	(8.9)	(50.0)	(18.5)	(18.5)	(1.6)			(0.8)	(100.0)				
合計	92	2,281	2,477	26,428	3,847	4,031	218	284	17	36	39,711	224	39,935	4.10	0.50
(%)	(0.2)	(5.7)	(6.2)	(66.6)	(9.7)	(10.2)	(0.5)	(0.7)	(0.0)	(0.1)	(100.0)				
不明		1	2	6	1	1					11		11	3.95	0.52
(%)		(9.1)	(18.2)	(54.5)	(9.1)	(9.1)					(100.0)				
記載なし															
(%)															
総計	92	2,282	2,479	26,434	3,848	4,032	218	284	17	36	39,722	224	39,946	4.10	0.50
(%)	(0.2)	(5.7)	(6.2)	(66.5)	(9.7)	(10.2)	(0.5)	(0.7)	(0.0)	(0.1)	(100.0)				
平均	5.43	4.39	7.25	9.85	12.81	13.07	13.83	13.38	17.12	11.58	10.03	9.23	10.02		
標準偏差	6.82	5.66	7.46	8.64	9.48	9.61	9.28	9.92	10.44	8.99	8.87	8.41	8.87		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査