

表 1929 透析時間(時間/回) 体重減少量(kg)別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

体重減少量	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
0.0未満	1	17	10	56	9	3		1			97		97	3.88	0.54
(%)	(1.0)	(17.5)	(10.3)	(57.7)	(9.3)	(3.1)		(1.0)			(100.0)				
0.0～	7	189	119	805	78	101	8	7	2	1	1,317	3	1,320	3.95	0.56
(%)	(0.5)	(14.4)	(9.0)	(61.1)	(5.9)	(7.7)	(0.6)	(0.5)	(0.2)	(0.1)	(100.0)				
1.0～	24	597	587	4,962	549	596	35	38	2	5	7,395	5	7,400	4.03	0.50
(%)	(0.3)	(8.1)	(7.9)	(67.1)	(7.4)	(8.1)	(0.5)	(0.5)	(0.0)	(0.1)	(100.0)				
2.0～	28	483	920	10,127	1,436	1,444	77	99	5	11	14,630	24	14,654	4.12	0.46
(%)	(0.2)	(3.3)	(6.3)	(69.2)	(9.8)	(9.9)	(0.5)	(0.7)	(0.0)	(0.1)	(100.0)				
3.0～	2	112	255	5,469	1,043	1,000	46	65	4	10	8,006	8	8,014	4.21	0.46
(%)	(0.0)	(1.4)	(3.2)	(68.3)	(13.0)	(12.5)	(0.6)	(0.8)	(0.0)	(0.1)	(100.0)				
4.0～		14	56	1,197	371	482	34	33	1	2	2,190	1	2,191	4.36	0.51
(%)		(0.6)	(2.6)	(54.7)	(16.9)	(22.0)	(1.6)	(1.5)	(0.0)	(0.1)	(100.0)				
5.0～		4	6	156	77	129	9	16	1	2	400		400	4.56	0.63
(%)		(1.0)	(1.5)	(39.0)	(19.3)	(32.3)	(2.3)	(4.0)	(0.3)	(0.5)	(100.0)				
6.0～			1	20	7	24	1	3	1	2	59		59	4.75	0.78
(%)			(1.7)	(33.9)	(11.9)	(40.7)	(1.7)	(5.1)	(1.7)	(3.4)	(100.0)				
7.0～				4	2	7		2		1	16		16	4.97	0.92
(%)				(25.0)	(12.5)	(43.8)		(12.5)		(6.3)	(100.0)				
8.0～		2	2	20	3	3					30		30	4.06	0.43
(%)		(6.7)	(6.7)	(66.7)	(10.0)	(10.0)					(100.0)				
合計	62	1,418	1,956	22,816	3,575	3,789	210	264	16	34	34,140	41	34,181	4.13	0.49
(%)	(0.2)	(4.2)	(5.7)	(66.8)	(10.5)	(11.1)	(0.6)	(0.8)	(0.0)	(0.1)	(100.0)				
記載なし	1	27	41	487	51	34	1	1			643	147	790	4.02	0.38
(%)	(0.2)	(4.2)	(6.4)	(75.7)	(7.9)	(5.3)	(0.2)	(0.2)			(100.0)				
総計	63	1,445	1,997	23,303	3,626	3,823	211	265	16	34	34,783	188	34,971	4.13	0.49
(%)	(0.2)	(4.2)	(5.7)	(67.0)	(10.4)	(11.0)	(0.6)	(0.8)	(0.0)	(0.1)	(100.0)				
平均	1.79	1.85	2.24	2.54	2.82	2.91	2.94	2.99	2.92	3.17	2.57	2.43	2.57		
標準偏差	0.83	0.95	1.00	0.98	1.03	1.12	1.17	1.26	1.53	1.59	1.03	0.81	1.03		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査