

表 1930 透析時間(時間/回) 体重減少率(%)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

体重減少率	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
1.0未満	6	84	48	366	40	38	2	2		1	587		587	3.93	0.55
(%)	(1.0)	(14.3)	(8.2)	(62.4)	(6.8)	(6.5)	(0.3)	(0.3)		(0.2)	(100.0)				
1.0～	4	127	98	674	70	96	7	9	2	1	1,088	3	1,091	4.00	0.56
(%)	(0.4)	(11.7)	(9.0)	(61.9)	(6.4)	(8.8)	(0.6)	(0.8)	(0.2)	(0.1)	(100.0)				
2.0～	10	235	199	1,995	273	275	13	22		4	3,026	4	3,030	4.06	0.52
(%)	(0.3)	(7.8)	(6.6)	(65.9)	(9.0)	(9.1)	(0.4)	(0.7)		(0.1)	(100.0)				
3.0～	16	364	444	4,606	612	744	40	50	5	5	6,886	7	6,893	4.11	0.50
(%)	(0.2)	(5.3)	(6.4)	(66.9)	(8.9)	(10.8)	(0.6)	(0.7)	(0.1)	(0.1)	(100.0)				
4.0～	16	321	535	6,357	1,009	986	60	71	3	7	9,365	12	9,377	4.14	0.47
(%)	(0.2)	(3.4)	(5.7)	(67.9)	(10.8)	(10.5)	(0.6)	(0.8)	(0.0)	(0.1)	(100.0)				
5.0～	8	169	364	5,019	863	873	53	56	2	6	7,413	11	7,424	4.17	0.47
(%)	(0.1)	(2.3)	(4.9)	(67.7)	(11.6)	(11.8)	(0.7)	(0.8)	(0.0)	(0.1)	(100.0)				
6.0～	1	76	173	2,511	423	482	16	35	2	4	3,723	2	3,725	4.19	0.47
(%)	(0.0)	(2.0)	(4.6)	(67.4)	(11.4)	(12.9)	(0.4)	(0.9)	(0.1)	(0.1)	(100.0)				
7.0～	1	25	57	915	191	191	13	10		3	1,406	2	1,408	4.22	0.48
(%)	(0.1)	(1.8)	(4.1)	(65.1)	(13.6)	(13.6)	(0.9)	(0.7)		(0.2)	(100.0)				
8.0～		7	24	257	67	67	6	5	1	2	436		436	4.27	0.55
(%)		(1.6)	(5.5)	(58.9)	(15.4)	(15.4)	(1.4)	(1.1)	(0.2)	(0.5)	(100.0)				
9.0～		8	12	101	26	37		4	1	1	190		190	4.28	0.65
(%)		(4.2)	(6.3)	(53.2)	(13.7)	(19.5)		(2.1)	(0.5)	(0.5)	(100.0)				
合計	62	1,416	1,954	22,801	3,574	3,789	210	264	16	34	34,120	41	34,161	4.13	0.49
(%)	(0.2)	(4.2)	(5.7)	(66.8)	(10.5)	(11.1)	(0.6)	(0.8)	(0.0)	(0.1)	(100.0)				
記載なし	1	29	43	502	52	34	1	1			663	147	810	4.02	0.38
(%)	(0.2)	(4.4)	(6.5)	(75.7)	(7.8)	(5.1)	(0.2)	(0.2)			(100.0)				
総計	63	1,445	1,997	23,303	3,626	3,823	211	265	16	34	34,783	188	34,971	4.13	0.49
(%)	(0.2)	(4.2)	(5.7)	(67.0)	(10.4)	(11.0)	(0.6)	(0.8)	(0.0)	(0.1)	(100.0)				
平均	3.56	3.73	4.33	4.60	4.83	4.80	4.73	4.74	4.90	4.97	4.59	4.48	4.59		
標準偏差	1.57	1.70	1.60	1.55	1.57	1.59	1.56	1.75	2.40	2.12	1.59	1.42	1.59		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査