

表 1945 透析時間(時間/回) 総コレステロール濃度(mg/dL)別 (血液透析濾過患者 透析回数週3回)

総コレステロール濃度	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
100未満	4	54	63	713	80	65	4	6	1		990	2	992	4.05	0.46
(%)	(0.4)	(5.5)	(6.4)	(72.0)	(8.1)	(6.6)	(0.4)	(0.6)	(0.1)		(100.0)				
100～	7	180	177	2,063	247	288	7	12		2	2,983	5	2,988	4.07	0.48
(%)	(0.2)	(6.0)	(5.9)	(69.2)	(8.3)	(9.7)	(0.2)	(0.4)		(0.1)	(100.0)				
120～	7	307	321	3,857	554	552	31	46	2	4	5,681	13	5,694	4.10	0.49
(%)	(0.1)	(5.4)	(5.7)	(67.9)	(9.8)	(9.7)	(0.5)	(0.8)	(0.0)	(0.1)	(100.0)				
140～	15	347	408	4,650	639	721	49	57	4	7	6,897	8	6,905	4.12	0.51
(%)	(0.2)	(5.0)	(5.9)	(67.4)	(9.3)	(10.5)	(0.7)	(0.8)	(0.1)	(0.1)	(100.0)				
160～	9	294	334	4,001	599	668	41	50	4	8	6,008	3	6,011	4.13	0.51
(%)	(0.1)	(4.9)	(5.6)	(66.6)	(10.0)	(11.1)	(0.7)	(0.8)	(0.1)	(0.1)	(100.0)				
180～	8	203	221	2,495	415	479	19	25		4	3,869	4	3,873	4.13	0.51
(%)	(0.2)	(5.2)	(5.7)	(64.5)	(10.7)	(12.4)	(0.5)	(0.6)		(0.1)	(100.0)				
200～	4	128	133	1,408	244	270	23	33	2	4	2,249	4	2,253	4.15	0.56
(%)	(0.2)	(5.7)	(5.9)	(62.6)	(10.8)	(12.0)	(1.0)	(1.5)	(0.1)	(0.2)	(100.0)				
220～		55	61	611	112	106	3	5		1	954	2	956	4.11	0.50
(%)		(5.8)	(6.4)	(64.0)	(11.7)	(11.1)	(0.3)	(0.5)		(0.1)	(100.0)				
240～		21	28	258	24	49	1	2			383	1	384	4.10	0.49
(%)		(5.5)	(7.3)	(67.4)	(6.3)	(12.8)	(0.3)	(0.5)			(100.0)				
260～	1	23	15	134	20	22	1				216		216	4.03	0.51
(%)	(0.5)	(10.6)	(6.9)	(62.0)	(9.3)	(10.2)	(0.5)				(100.0)				
合計	55	1,612	1,761	20,190	2,934	3,220	179	236	13	30	30,230	42	30,272	4.11	0.50
(%)	(0.2)	(5.3)	(5.8)	(66.8)	(9.7)	(10.7)	(0.6)	(0.8)	(0.0)	(0.1)	(100.0)				
記載なし	37	670	718	6,244	914	812	39	48	4	6	9,492	182	9,674	4.05	0.49
(%)	(0.4)	(7.1)	(7.6)	(65.8)	(9.6)	(8.6)	(0.4)	(0.5)	(0.0)	(0.1)	(100.0)				
総計	92	2,282	2,479	26,434	3,848	4,032	218	284	17	36	39,722	224	39,946	4.10	0.50
(%)	(0.2)	(5.7)	(6.2)	(66.5)	(9.7)	(10.2)	(0.5)	(0.7)	(0.0)	(0.1)	(100.0)				
平均	153.25	158.37	158.30	157.00	160.43	161.37	161.65	161.53	154.46	165.30	158.01	152.71	158.00		
標準偏差	35.75	37.92	37.06	35.61	35.11	34.90	32.73	32.86	32.62	28.27	35.69	39.44	35.70		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査