

表 1946 透析時間(時間/回) HDL-C濃度(mg/dL)別(血液透析濾過患者 透析回数週3回)

HDL-C濃度	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
20未満	2	10	4	51	9	4			1		81		81	3.97	0.61
(%)	(2.5)	(12.3)	(4.9)	(63.0)	(11.1)	(4.9)			(1.2)		(100.0)				
20～	3	106	100	1,228	186	219	14	15			1,871	3	1,874	4.12	0.50
(%)	(0.2)	(5.7)	(5.3)	(65.6)	(9.9)	(11.7)	(0.7)	(0.8)			(100.0)				
30～	11	330	371	4,114	619	601	34	35	1	5	6,121	10	6,131	4.10	0.49
(%)	(0.2)	(5.4)	(6.1)	(67.2)	(10.1)	(9.8)	(0.6)	(0.6)	(0.0)	(0.1)	(100.0)				
40～	10	380	463	4,936	722	734	42	55	3	6	7,351	9	7,360	4.11	0.49
(%)	(0.1)	(5.2)	(6.3)	(67.1)	(9.8)	(10.0)	(0.6)	(0.7)	(0.0)	(0.1)	(100.0)				
50～	9	289	350	4,036	582	582	43	59	4	7	5,961	10	5,971	4.12	0.50
(%)	(0.2)	(4.8)	(5.9)	(67.7)	(9.8)	(9.8)	(0.7)	(1.0)	(0.1)	(0.1)	(100.0)				
60～	2	169	256	2,547	412	401	24	34	2	8	3,855	2	3,857	4.13	0.50
(%)	(0.1)	(4.4)	(6.6)	(66.1)	(10.7)	(10.4)	(0.6)	(0.9)	(0.1)	(0.2)	(100.0)				
70～	2	95	134	1,332	200	228	7	16	2	4	2,020	3	2,023	4.13	0.51
(%)	(0.1)	(4.7)	(6.6)	(65.9)	(9.9)	(11.3)	(0.3)	(0.8)	(0.1)	(0.2)	(100.0)				
80～	3	33	60	669	103	133	6	4		1	1,012	1	1,013	4.15	0.47
(%)	(0.3)	(3.3)	(5.9)	(66.1)	(10.2)	(13.1)	(0.6)	(0.4)		(0.1)	(100.0)				
90～	4	24	25	276	38	56	2	5			430	1	431	4.12	0.54
(%)	(0.9)	(5.6)	(5.8)	(64.2)	(8.8)	(13.0)	(0.5)	(1.2)			(100.0)				
100～	1	20	15	255	37	45	3	5	1	1	383		383	4.18	0.60
(%)	(0.3)	(5.2)	(3.9)	(66.6)	(9.7)	(11.7)	(0.8)	(1.3)	(0.3)	(0.3)	(100.0)				
合計	47	1,456	1,778	19,444	2,908	3,003	175	228	14	32	29,085	39	29,124	4.12	0.50
(%)	(0.2)	(5.0)	(6.1)	(66.9)	(10.0)	(10.3)	(0.6)	(0.8)	(0.0)	(0.1)	(100.0)				
記載なし	45	826	701	6,990	940	1,029	43	56	3	4	10,637	185	10,822	4.05	0.50
(%)	(0.4)	(7.8)	(6.6)	(65.7)	(8.8)	(9.7)	(0.4)	(0.5)	(0.0)	(0.0)	(100.0)				
総計	92	2,282	2,479	26,434	3,848	4,032	218	284	17	36	39,722	224	39,946	4.10	0.50
(%)	(0.2)	(5.7)	(6.2)	(66.5)	(9.7)	(10.2)	(0.5)	(0.7)	(0.0)	(0.1)	(100.0)				
平均	52.11	49.54	50.89	50.79	50.95	51.63	50.33	52.59	55.57	56.94	50.86	48.56	50.85		
標準偏差	21.59	17.10	16.37	17.12	17.09	18.10	16.76	17.27	20.00	16.29	17.18	15.69	17.18		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査