

表 1959 血流量(ml/分) 透析後体重(kg)別（施設血液透析患者 透析回数週3回）

透析後体重	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差
20未満			1	3	3	11	50	7	6		2	1			84		84	202.02	27.45
(%)			(1.2)	(3.6)	(3.6)	(13.1)	(59.5)	(8.3)	(7.1)		(2.4)	(1.2)			(100.0)				
20～	1	14	65	250	54	147	225	24	10		1	1	2	2	796	6	802	172.50	35.33
(%)	(0.1)	(1.8)	(8.2)	(31.4)	(6.8)	(18.5)	(28.3)	(3.0)	(1.3)		(0.1)	(0.1)	(0.3)	(0.3)	(100.0)				
30～	9	130	496	3,818	1,224	4,111	7,425	878	609	41	24	63	14	51	18,893	224	19,117	184.54	32.74
(%)	(0.0)	(0.7)	(2.6)	(20.2)	(6.5)	(21.8)	(39.3)	(4.6)	(3.2)	(0.2)	(0.1)	(0.3)	(0.1)	(0.3)	(100.0)				
40～	14	223	710	6,697	2,316	10,927	28,422	5,215	3,539	272	187	428	40	150	59,140	669	59,809	195.02	32.06
(%)	(0.0)	(0.4)	(1.2)	(11.3)	(3.9)	(18.5)	(48.1)	(8.8)	(6.0)	(0.5)	(0.3)	(0.7)	(0.1)	(0.3)	(100.0)				
50～	13	199	478	4,490	1,488	9,396	34,701	9,940	8,066	674	600	991	107	170	71,313	796	72,109	205.08	32.92
(%)	(0.0)	(0.3)	(0.7)	(6.3)	(2.1)	(13.2)	(48.7)	(13.9)	(11.3)	(0.9)	(0.8)	(1.4)	(0.2)	(0.2)	(100.0)				
60～	2	101	216	1,600	551	3,830	18,987	7,888	7,687	781	639	1,125	84	128	43,619	463	44,082	214.28	34.87
(%)	(0.0)	(0.2)	(0.5)	(3.7)	(1.3)	(8.8)	(43.5)	(18.1)	(17.6)	(1.8)	(1.5)	(2.6)	(0.2)	(0.3)	(100.0)				
70～	2	28	77	485	138	1,042	5,987	3,438	3,976	434	483	770	79	54	16,993	191	17,184	222.90	37.48
(%)	(0.0)	(0.2)	(0.5)	(2.9)	(0.8)	(6.1)	(35.2)	(20.2)	(23.4)	(2.6)	(2.8)	(4.5)	(0.5)	(0.3)	(100.0)				
80～	3	11	25	132	37	302	1,748	1,160	1,679	190	237	425	45	26	6,020	61	6,081	229.95	39.49
(%)	(0.0)	(0.2)	(0.4)	(2.2)	(0.6)	(5.0)	(29.0)	(19.3)	(27.9)	(3.2)	(3.9)	(7.1)	(0.7)	(0.4)	(100.0)				
90～		6	9	43	6	81	557	378	671	76	95	216	27	13	2,178	17	2,195	236.07	42.03
(%)		(0.3)	(0.4)	(2.0)	(0.3)	(3.7)	(25.6)	(17.4)	(30.8)	(3.5)	(4.4)	(9.9)	(1.2)	(0.6)	(100.0)				
100～		4	5	12	17	160	112	254	23	38	105	13	4	747	8	755	242.12	43.32	
(%)		(0.5)	(0.7)	(1.6)	(2.3)	(21.4)	(15.0)	(34.0)	(3.1)	(5.1)	(14.1)	(1.7)	(0.5)	(100.0)					
110～			3	6	1	9	57	38	89	15	12	56	7	3	296	3	299	247.56	49.22
(%)			(1.0)	(2.0)	(0.3)	(3.0)	(19.3)	(12.8)	(30.1)	(5.1)	(4.1)	(18.9)	(2.4)	(1.0)	(100.0)				
120～				5		5	31	20	58	14	9	36	8	2	188	1	189	252.82	49.16
(%)				(2.7)		(2.7)	(16.5)	(10.6)	(30.9)	(7.4)	(4.8)	(19.1)	(4.3)	(1.1)	(100.0)				
合計	44	716	2,085	17,541	5,818	29,878	98,350	29,098	26,644	2,520	2,327	4,217	426	603	220,267	2,439	222,706	204.91	35.89
(%)	(0.0)	(0.3)	(0.9)	(8.0)	(2.6)	(13.6)	(44.7)	(13.2)	(12.1)	(1.1)	(1.1)	(1.9)	(0.2)	(0.3)	(100.0)				
記載なし		23	63	432	83	607	2,140	448	416	13	10	48	3	115	4,401	1,864	6,265	206.50	56.76
(%)		(0.5)	(1.4)	(9.8)	(1.9)	(13.8)	(48.6)	(10.2)	(9.5)	(0.3)	(0.2)	(1.1)	(0.1)	(2.6)	(100.0)				
総計	44	739	2,148	17,973	5,901	30,485	100,490	29,546	27,060	2,533	2,337	4,265	429	718	224,668	4,303	228,971	204.94	36.42
(%)	(0.0)	(0.3)	(1.0)	(8.0)	(2.6)	(13.6)	(44.7)	(13.2)	(12.0)	(1.1)	(1.0)	(1.9)	(0.2)	(0.3)	(100.0)				
平均	50.40	50.76	48.29	48.16	48.10	50.83	54.47	59.61	62.71	64.77	66.86	67.73	68.71	57.67	55.45	54.87	55.45		
標準偏差	13.91	12.89	12.57	11.10	10.36	10.72	11.42	12.28	13.75	14.47	14.63	16.72	19.46	14.87	12.90	12.50	12.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査