

表 1963 血流量(ml/分) 透析前BUN(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前BUN	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差
20未満	2	4	10	70	17	78	214	33	34	2	1	4			469	10	479	191.29	31.97
(%)	(0.4)	(0.9)	(2.1)	(14.9)	(3.6)	(16.6)	(45.6)	(7.0)	(7.2)	(0.4)	(0.2)	(0.9)			(100.0)				
20～		10	44	304	79	364	1,148	226	196	12	13	34	6	7	2,443	37	2,480	197.21	35.25
(%)		(0.4)	(1.8)	(12.4)	(3.2)	(14.9)	(47.0)	(9.3)	(8.0)	(0.5)	(0.5)	(1.4)	(0.2)	(0.3)	(100.0)				
30～	3	23	104	944	313	1,551	4,574	1,193	1,109	106	95	194	14	18	10,241	117	10,358	202.29	34.69
(%)	(0.0)	(0.2)	(1.0)	(9.2)	(3.1)	(15.1)	(44.7)	(11.6)	(10.8)	(1.0)	(0.9)	(1.9)	(0.1)	(0.2)	(100.0)				
40～	2	55	178	1,756	737	3,622	12,026	3,485	3,277	340	274	568	65	92	26,477	334	26,811	206.55	36.15
(%)	(0.0)	(0.2)	(0.7)	(6.6)	(2.8)	(13.7)	(45.4)	(13.2)	(12.4)	(1.3)	(1.0)	(2.1)	(0.2)	(0.3)	(100.0)				
50～	3	44	226	2,376	981	5,388	19,024	6,140	5,682	592	545	918	107	122	42,148	511	42,659	208.65	35.13
(%)	(0.0)	(0.1)	(0.5)	(5.6)	(2.3)	(12.8)	(45.1)	(14.6)	(13.5)	(1.4)	(1.3)	(2.2)	(0.3)	(0.3)	(100.0)				
60～	1	41	192	2,137	991	5,131	18,959	6,441	5,906	579	553	954	87	148	42,120	523	42,643	209.74	35.44
(%)	(0.0)	(0.1)	(0.5)	(5.1)	(2.4)	(12.2)	(45.0)	(15.3)	(14.0)	(1.4)	(1.3)	(2.3)	(0.2)	(0.4)	(100.0)				
70～	1	31	143	1,362	663	3,272	12,378	4,488	4,099	405	366	639	69	68	27,984	338	28,322	210.22	34.72
(%)	(0.0)	(0.1)	(0.5)	(4.9)	(2.4)	(11.7)	(44.2)	(16.0)	(14.6)	(1.4)	(1.3)	(2.3)	(0.2)	(0.2)	(100.0)				
80～		21	50	638	345	1,531	5,741	2,171	1,890	175	165	269	15	24	13,035	177	13,212	209.58	33.46
(%)		(0.2)	(0.4)	(4.9)	(2.6)	(11.7)	(44.0)	(16.7)	(14.5)	(1.3)	(1.3)	(2.1)	(0.1)	(0.2)	(100.0)				
90～	2	5	34	269	122	547	2,091	743	691	59	58	103	8	8	4,740	63	4,803	208.90	34.22
(%)	(0.0)	(0.1)	(0.7)	(5.7)	(2.6)	(11.5)	(44.1)	(15.7)	(14.6)	(1.2)	(1.2)	(2.2)	(0.2)	(0.2)	(100.0)				
100～		5	7	55	42	154	566	222	198	23	21	31	5		1,329	19	1,348	210.36	33.31
(%)		(0.4)	(0.5)	(4.1)	(3.2)	(11.6)	(42.6)	(16.7)	(14.9)	(1.7)	(1.6)	(2.3)	(0.4)		(100.0)				
110～		4	1	21	7	43	154	65	61	6	8	13	1	2	386	8	394	213.06	41.29
(%)		(1.0)	(0.3)	(5.4)	(1.8)	(11.1)	(39.9)	(16.8)	(15.8)	(1.6)	(2.1)	(3.4)	(0.3)	(0.5)	(100.0)				
120～		3	2	18	6	26	91	25	19	3	3	7			203	1	204	202.66	36.48
(%)		(1.5)	(1.0)	(8.9)	(3.0)	(12.8)	(44.8)	(12.3)	(9.4)	(1.5)	(1.5)	(3.4)			(100.0)				
合計	14	246	991	9,950	4,303	21,707	76,966	25,232	23,162	2,302	2,102	3,734	377	489	171,575	2,138	173,713	208.35	35.21
(%)	(0.0)	(0.1)	(0.6)	(5.8)	(2.5)	(12.7)	(44.9)	(14.7)	(13.5)	(1.3)	(1.2)	(2.2)	(0.2)	(0.3)	(100.0)				
記載なし		1	10	198	68	429	1,652	379	323	16	13	47	8	101	3,245	1,251	4,496	212.73	58.58
(%)		(0.0)	(0.3)	(6.1)	(2.1)	(13.2)	(50.9)	(11.7)	(10.0)	(0.5)	(0.4)	(1.4)	(0.2)	(3.1)	(100.0)				
総計	14	247	1,001	10,148	4,371	22,136	78,618	25,611	23,485	2,318	2,115	3,781	385	590	174,820	3,389	178,209	208.43	35.79
(%)	(0.0)	(0.1)	(0.6)	(5.8)	(2.5)	(12.7)	(45.0)	(14.6)	(13.4)	(1.3)	(1.2)	(2.2)	(0.2)	(0.3)	(100.0)				
平均	51.21	59.00	57.28	58.03	60.25	59.86	60.76	62.39	62.20	62.06	62.51	61.61	60.22	59.69	60.94	61.11	60.94		
標準偏差	23.69	21.29	17.91	16.84	16.48	15.82	15.58	15.21	15.15	14.84	14.93	15.30	14.66	13.66	15.63	16.10	15.64		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査