

表 1964 血流量(ml/分) 透析前クレアチニン濃度(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前クレアチニン濃度	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差
2未満		1	8	22	7	7	22	7	1						75	1	76	173.33	32.56
(%)		(1.3)	(10.7)	(29.3)	(9.3)	(9.3)	(29.3)	(9.3)	(1.3)						(100.0)				
2～	1	12	74	420	101	303	574	77	53	2			8	1	1,628	28	1,656	180.37	32.64
(%)	(0.1)	(0.7)	(4.5)	(25.8)	(6.2)	(18.6)	(35.3)	(4.7)	(3.3)	(0.1)			(0.5)	(0.1)	(100.0)				
4～	3	51	155	1,463	363	1,488	3,218	482	343	16	14		54	6	7,678	117	7,795	188.21	34.35
(%)	(0.0)	(0.7)	(2.0)	(19.1)	(4.7)	(19.4)	(41.9)	(6.3)	(4.5)	(0.2)	(0.2)		(0.7)	(0.1)	(100.0)				
6～	4	59	241	2,632	939	4,447	10,944	2,124	1,708	136	102		236	32	23,668	338	24,006	196.50	32.97
(%)	(0.0)	(0.2)	(1.0)	(11.1)	(4.0)	(18.8)	(46.2)	(9.0)	(7.2)	(0.6)	(0.4)		(1.0)	(0.1)	(100.0)				
8～	1	59	278	2,877	1,416	6,989	22,192	5,506	4,589	453	361		676	77	45,584	531	46,115	203.56	32.73
(%)	(0.0)	(0.1)	(0.6)	(6.3)	(3.1)	(15.3)	(48.7)	(12.1)	(10.1)	(1.0)	(0.8)		(1.5)	(0.2)	(100.0)				
10～	4	37	154	1,688	957	5,392	22,280	7,671	6,702	707	602	1,046	105	142	47,487	547	48,034	211.02	33.79
(%)	(0.0)	(0.1)	(0.3)	(3.6)	(2.0)	(11.4)	(46.9)	(16.2)	(14.1)	(1.5)	(1.3)		(2.2)	(0.2)	(100.0)				
12～		15	60	636	397	2,288	12,185	5,827	5,698	600	589		954	100	29,441	313	29,754	218.42	34.74
(%)		(0.1)	(0.2)	(2.2)	(1.3)	(7.8)	(41.4)	(19.8)	(19.4)	(2.0)	(2.0)		(3.2)	(0.3)	(100.0)				
14～	1	10	17	167	90	634	4,220	2,622	2,964	277	317		537	36	11,929	107	12,036	224.73	35.21
(%)	(0.0)	(0.1)	(0.1)	(1.4)	(0.8)	(5.3)	(35.4)	(22.0)	(24.8)	(2.3)	(2.7)		(4.5)	(0.3)	(100.0)				
16～		2	4	43	29	150	1,323	911	1,104	111	117		221	20	4,054	46	4,100	228.80	37.34
(%)		(0.0)	(0.1)	(1.1)	(0.7)	(3.7)	(32.6)	(22.5)	(27.2)	(2.7)	(2.9)		(5.5)	(0.5)	(100.0)				
合計	14	246	991	9,948	4,299	21,698	76,958	25,227	23,162	2,302	2,102	3,732	377	488	171,544	2,028	173,572	208.35	35.21
(%)	(0.0)	(0.1)	(0.6)	(5.8)	(2.5)	(12.6)	(44.9)	(14.7)	(13.5)	(1.3)	(1.2)		(2.2)	(0.2)	(100.0)				
記載なし		1	10	200	72	438	1,660	384	323	16	13		49	8	3,276	1,361	4,637	212.67	58.51
(%)		(0.0)	(0.3)	(6.1)	(2.2)	(13.4)	(50.7)	(11.7)	(9.9)	(0.5)	(0.4)		(1.5)	(0.2)	(100.0)				
総計	14	247	1,001	10,148	4,371	22,136	78,618	25,611	23,485	2,318	2,115	3,781	385	590	174,820	3,389	178,209	208.43	35.79
(%)	(0.0)	(0.1)	(0.6)	(5.8)	(2.5)	(12.7)	(45.0)	(14.6)	(13.4)	(1.3)	(1.2)		(2.2)	(0.2)	(100.0)				
平均	7.90	8.15	8.07	8.30	9.02	9.34	10.14	11.15	11.47	11.60	11.89	11.71	11.29	10.65	10.30	9.95	10.29		
標準偏差	3.21	2.98	3.00	2.66	2.53	2.55	2.63	2.63	2.69	2.64	2.55	2.70	2.70	2.93	2.79	2.79	2.79		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査