

表 1969 血流量(ml/分) 透析前アルブミン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前アルブミン濃度	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差
1.5未満		6	16	37	4	20	51	4	4	1				1	144	2	146	174.93	44.72
(%)		(4.2)	(11.1)	(25.7)	(2.8)	(13.9)	(35.4)	(2.8)	(2.8)	(0.7)				(0.7)	(100.0)				
1.5～	1	21	59	204	32	114	272	35	16	1	1	2		2	760	11	771	176.31	36.49
(%)	(0.1)	(2.8)	(7.8)	(26.8)	(4.2)	(15.0)	(35.8)	(4.6)	(2.1)	(0.1)	(0.1)	(0.3)		(0.3)	(100.0)				
2.0～	8	66	130	789	127	524	1,194	142	122	6	1	10		6	3,125	54	3,179	180.81	34.84
(%)	(0.3)	(2.1)	(4.2)	(25.2)	(4.1)	(16.8)	(38.2)	(4.5)	(3.9)	(0.2)	(0.0)	(0.3)		(0.2)	(100.0)				
2.5～	11	138	312	2,130	495	2,151	5,058	823	691	41	31	75	4	28	11,988	162	12,150	188.93	34.92
(%)	(0.1)	(1.2)	(2.6)	(17.8)	(4.1)	(17.9)	(42.2)	(6.9)	(5.8)	(0.3)	(0.3)	(0.6)	(0.0)	(0.2)	(100.0)				
3.0～	10	226	659	5,557	1,678	8,174	23,340	5,461	4,793	357	367	645	73	106	51,446	636	52,082	198.96	34.47
(%)	(0.0)	(0.4)	(1.3)	(10.8)	(3.3)	(15.9)	(45.4)	(10.6)	(9.3)	(0.7)	(0.7)	(1.3)	(0.1)	(0.2)	(100.0)				
3.5～	9	182	693	6,577	2,501	13,680	48,157	15,075	13,663	1,403	1,239	2,240	202	313	105,934	1,201	107,135	207.52	35.30
(%)	(0.0)	(0.2)	(0.7)	(6.2)	(2.4)	(12.9)	(45.5)	(14.2)	(12.9)	(1.3)	(1.2)	(2.1)	(0.2)	(0.3)	(100.0)				
4.0～	1	53	145	1,675	721	4,202	17,465	6,688	6,501	623	620	1,125	126	136	40,081	425	40,506	213.13	35.82
(%)	(0.0)	(0.1)	(0.4)	(4.2)	(1.8)	(10.5)	(43.6)	(16.7)	(16.2)	(1.6)	(1.5)	(2.8)	(0.3)	(0.3)	(100.0)				
4.5～		2	20	86	29	203	1,020	394	426	39	40	66	7	10	2,342	20	2,362	215.05	36.94
(%)		(0.1)	(0.9)	(3.7)	(1.2)	(8.7)	(43.6)	(16.8)	(18.2)	(1.7)	(1.7)	(2.8)	(0.3)	(0.4)	(100.0)				
5.0～		2	2	13	3	27	85	22	11	2			4		171	1	172	199.65	31.42
(%)		(1.2)	(1.2)	(7.6)	(1.8)	(15.8)	(49.7)	(12.9)	(6.4)	(1.2)			(2.3)		(100.0)				
合計	40	696	2,036	17,068	5,590	29,095	96,642	28,644	26,227	2,473	2,299	4,167	412	602	215,991	2,512	218,503	205.05	35.92
(%)	(0.0)	(0.3)	(0.9)	(7.9)	(2.6)	(13.5)	(44.7)	(13.3)	(12.1)	(1.1)	(1.1)	(1.9)	(0.2)	(0.3)	(100.0)				
記載なし	4	43	112	905	311	1,390	3,848	902	833	60	38	98	17	116	8,677	1,791	10,468	202.17	47.03
(%)	(0.0)	(0.5)	(1.3)	(10.4)	(3.6)	(16.0)	(44.3)	(10.4)	(9.6)	(0.7)	(0.4)	(1.1)	(0.2)	(1.3)	(100.0)				
総計	44	739	2,148	17,973	5,901	30,485	100,490	29,546	27,060	2,533	2,337	4,265	429	718	224,668	4,303	228,971	204.94	36.42
(%)	(0.0)	(0.3)	(1.0)	(8.0)	(2.6)	(13.6)	(44.7)	(13.2)	(12.0)	(1.1)	(1.0)	(1.9)	(0.2)	(0.3)	(100.0)				
平均	3.00	3.17	3.26	3.37	3.48	3.53	3.60	3.70	3.72	3.75	3.76	3.75	3.76	3.67	3.60	3.56	3.60		
標準偏差	0.58	0.64	0.60	0.52	0.46	0.45	0.43	0.39	0.38	0.35	0.35	0.36	0.35	0.44	0.44	0.46	0.44		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査