

表 1972 血流量(ml/分) 透析前リン濃度(mg/dL)別 (施設血液透析患者 透析回数週3回)

透析前リン濃度	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差
3未満	6	63	172	1,300	301	1,449	3,905	782	655	42	29	93	10	21	8,828	133	8,961	194.02	35.29
(%)	(0.1)	(0.7)	(1.9)	(14.7)	(3.4)	(16.4)	(44.2)	(8.9)	(7.4)	(0.5)	(0.3)	(1.1)	(0.1)	(0.2)	(100.0)				
3.0～	8	64	153	1,248	351	1,703	4,923	1,116	933	77	57	161	13	32	10,839	141	10,980	198.18	35.92
(%)	(0.1)	(0.6)	(1.4)	(11.5)	(3.2)	(15.7)	(45.4)	(10.3)	(8.6)	(0.7)	(0.5)	(1.5)	(0.1)	(0.3)	(100.0)				
3.5～	5	75	254	2,001	536	2,737	8,498	2,110	1,880	160	163	298	39	51	18,807	211	19,018	200.77	36.29
(%)	(0.0)	(0.4)	(1.4)	(10.6)	(2.9)	(14.6)	(45.2)	(11.2)	(10.0)	(0.9)	(0.9)	(1.6)	(0.2)	(0.3)	(100.0)				
4.0～	3	89	307	2,535	800	4,106	12,330	3,416	3,172	258	269	484	53	52	27,874	318	28,192	202.67	34.93
(%)	(0.0)	(0.3)	(1.1)	(9.1)	(2.9)	(14.7)	(44.2)	(12.3)	(11.4)	(0.9)	(1.0)	(1.7)	(0.2)	(0.2)	(100.0)				
4.5～	3	89	274	2,518	867	4,571	15,053	4,276	3,928	386	343	611	75	68	33,062	406	33,468	204.79	34.65
(%)	(0.0)	(0.3)	(0.8)	(7.6)	(2.6)	(13.8)	(45.5)	(12.9)	(11.9)	(1.2)	(1.0)	(1.8)	(0.2)	(0.2)	(100.0)				
5.0～	3	90	268	2,335	894	4,491	15,116	4,547	4,162	373	360	642	68	88	33,437	372	33,809	205.88	35.37
(%)	(0.0)	(0.3)	(0.8)	(7.0)	(2.7)	(13.4)	(45.2)	(13.6)	(12.4)	(1.1)	(1.1)	(1.9)	(0.2)	(0.3)	(100.0)				
5.5～		66	215	1,868	682	3,667	12,869	4,099	3,801	384	334	584	51	87	28,707	305	29,012	207.36	35.51
(%)		(0.2)	(0.7)	(6.5)	(2.4)	(12.8)	(44.8)	(14.3)	(13.2)	(1.3)	(1.2)	(2.0)	(0.2)	(0.3)	(100.0)				
6.0～	1	16	35	306	115	631	2,175	705	626	76	74	93	7	16	4,876	52	4,928	207.52	35.87
(%)	(0.0)	(0.3)	(0.7)	(6.3)	(2.4)	(12.9)	(44.6)	(14.5)	(12.8)	(1.6)	(1.5)	(1.9)	(0.1)	(0.3)	(100.0)				
6.1～	1	42	117	1,050	389	2,024	7,215	2,359	2,236	223	199	378	34	56	16,323	182	16,505	208.23	36.49
(%)	(0.0)	(0.3)	(0.7)	(6.4)	(2.4)	(12.4)	(44.2)	(14.5)	(13.7)	(1.4)	(1.2)	(2.3)	(0.2)	(0.3)	(100.0)				
6.5～	2	39	99	899	336	1,737	6,207	2,181	1,930	226	171	308	26	44	14,205	172	14,377	208.33	35.99
(%)	(0.0)	(0.3)	(0.7)	(6.3)	(2.4)	(12.2)	(43.7)	(15.4)	(13.6)	(1.6)	(1.2)	(2.2)	(0.2)	(0.3)	(100.0)				
7.0～	1	22	54	536	207	1,063	3,828	1,351	1,233	111	125	176	21	32	8,760	100	8,860	208.88	36.66
(%)	(0.0)	(0.3)	(0.6)	(6.1)	(2.4)	(12.1)	(43.7)	(15.4)	(14.1)	(1.3)	(1.4)	(2.0)	(0.2)	(0.4)	(100.0)				
7.5～	2	11	32	312	111	659	2,318	829	744	69	70	131	7	20	5,315	44	5,359	209.28	36.37
(%)	(0.0)	(0.2)	(0.6)	(5.9)	(2.1)	(12.4)	(43.6)	(15.6)	(14.0)	(1.3)	(1.3)	(2.5)	(0.1)	(0.4)	(100.0)				
8.0～	2	8	32	184	45	342	1,352	493	495	61	49	81	3	15	3,162	30	3,192	211.18	37.98
(%)	(0.1)	(0.3)	(1.0)	(5.8)	(1.4)	(10.8)	(42.8)	(15.6)	(15.7)	(1.9)	(1.5)	(2.6)	(0.1)	(0.5)	(100.0)				
8.5～	1	7	13	132	46	204	904	283	278	27	38	60	5	6	2,004	19	2,023	209.91	38.20
(%)	(0.0)	(0.3)	(0.6)	(6.6)	(2.3)	(10.2)	(45.1)	(14.1)	(13.9)	(1.3)	(1.9)	(3.0)	(0.2)	(0.3)	(100.0)				
9.0～	1	18	25	190	71	363	1,291	477	474	40	41	107	11	14	3,123	33	3,156	210.70	39.69
(%)	(0.0)	(0.6)	(0.8)	(6.1)	(2.3)	(11.6)	(41.3)	(15.3)	(15.2)	(1.3)	(1.3)	(3.4)	(0.4)	(0.4)	(100.0)				
合計	39	699	2,050	17,414	5,751	29,747	97,984	29,024	26,547	2,513	2,322	4,207	423	602	219,322	2,518	221,840	204.96	35.87
(%)	(0.0)	(0.3)	(0.9)	(7.9)	(2.6)	(13.6)	(44.7)	(13.2)	(12.1)	(1.1)	(1.1)	(1.9)	(0.2)	(0.3)	(100.0)				
記載なし	5	40	98	559	150	738	2,506	522	513	20	15	58	6	116	5,346	1,785	7,131	203.96	54.31
(%)	(0.1)	(0.7)	(1.8)	(10.5)	(2.8)	(13.8)	(46.9)	(9.8)	(9.6)	(0.4)	(0.3)	(1.1)	(0.1)	(2.2)	(100.0)				
総計	44	739	2,148	17,973	5,901	30,485	100,490	29,546	27,060	2,533	2,337	4,265	429	718	224,668	4,303	228,971	204.94	36.42
(%)	(0.0)	(0.3)	(1.0)	(8.0)	(2.6)	(13.6)	(44.7)	(13.2)	(12.0)	(1.1)	(1.0)	(1.9)	(0.2)	(0.3)	(100.0)				
平均	4.57	4.91	4.83	4.87	5.04	5.07	5.18	5.36	5.39	5.49	5.51	5.45	5.30	5.44	5.20	5.11	5.20		
標準偏差	2.19	1.67	1.55	1.47	1.43	1.41	1.41	1.40	1.41	1.38	1.40	1.49	1.45	1.51	1.43	1.44	1.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査