

表 1974 血流量(ml/分) PTH値(pg/mL)別(施設血液透析患者 透析回数週3回) PTH測定法:whole-PTH

PTH値	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差
20未満		2	8	128	45	256	646	163	166	18	15	30	2	1	1,480	19	1,499	202.83	33.15
(%)		(0.1)	(0.5)	(8.6)	(3.0)	(17.3)	(43.6)	(11.0)	(11.2)	(1.2)	(1.0)	(2.0)	(0.1)	(0.1)	(100.0)				
20～		2	20	120	52	263	837	239	220	22	17	42	4	2	1,840	8	1,848	205.27	33.23
(%)		(0.1)	(1.1)	(6.5)	(2.8)	(14.3)	(45.5)	(13.0)	(12.0)	(1.2)	(0.9)	(2.3)	(0.2)	(0.1)	(100.0)				
35～		2	10	110	53	170	712	237	210	22	14	55	2	3	1,600	13	1,613	208.18	34.74
(%)		(0.1)	(0.6)	(6.9)	(3.3)	(10.6)	(44.5)	(14.8)	(13.1)	(1.4)	(0.9)	(3.4)	(0.1)	(0.2)	(100.0)				
50～		2	19	144	63	289	1,085	311	358	31	39	96	9	1	2,447	12	2,459	210.05	35.69
(%)		(0.1)	(0.8)	(5.9)	(2.6)	(11.8)	(44.3)	(12.7)	(14.6)	(1.3)	(1.6)	(3.9)	(0.4)	(0.0)	(100.0)				
75～		2	14	113	39	225	914	285	357	32	19	127	4	4	2,135	13	2,148	213.67	37.11
(%)		(0.1)	(0.7)	(5.3)	(1.8)	(10.5)	(42.8)	(13.3)	(16.7)	(1.5)	(0.9)	(5.9)	(0.2)	(0.2)	(100.0)				
100～		2	17	121	36	198	732	259	261	28	26	86	6	2	1,774	13	1,787	211.31	37.48
(%)		(0.1)	(1.0)	(6.8)	(2.0)	(11.2)	(41.3)	(14.6)	(14.7)	(1.6)	(1.5)	(4.8)	(0.3)	(0.1)	(100.0)				
125～		1	21	79	29	108	488	180	176	22	16	55	3		1,178	9	1,187	210.87	37.32
(%)		(0.1)	(1.8)	(6.7)	(2.5)	(9.2)	(41.4)	(15.3)	(14.9)	(1.9)	(1.4)	(4.7)	(0.3)		(100.0)				
150～			1	6		1	10	8	10			1			37		37	211.08	39.98
(%)			(2.7)	(16.2)		(2.7)	(27.0)	(21.6)	(27.0)			(2.7)			(100.0)				
151～		1	12	63	21	93	392	137	134	20	15	40	1		929	5	934	210.52	36.19
(%)		(0.1)	(1.3)	(6.8)	(2.3)	(10.0)	(42.2)	(14.7)	(14.4)	(2.2)	(1.6)	(4.3)	(0.1)		(100.0)				
180～		2	7	47	21	74	271	91	105	11	7	24	1		661	5	666	208.78	35.82
(%)		(0.3)	(1.1)	(7.1)	(3.2)	(11.2)	(41.0)	(13.8)	(15.9)	(1.7)	(1.1)	(3.6)	(0.2)		(100.0)				
210～		3	7	29	19	32	210	68	69	5	5	18			465	7	472	208.67	35.74
(%)		(0.6)	(1.5)	(6.2)	(4.1)	(6.9)	(45.2)	(14.6)	(14.8)	(1.1)	(1.1)	(3.9)			(100.0)				
240～		2	7	41	16	64	218	78	60	11	4	18		2	521	5	526	206.87	38.45
(%)		(0.4)	(1.3)	(7.9)	(3.1)	(12.3)	(41.8)	(15.0)	(11.5)	(2.1)	(0.8)	(3.5)		(0.4)	(100.0)				
300～			1	16	8	22	70	36	28	5	4	11			201	4	205	212.09	37.90
(%)			(0.5)	(8.0)	(4.0)	(10.9)	(34.8)	(17.9)	(13.9)	(2.5)	(2.0)	(5.5)			(100.0)				
350～			4	6	3	13	63	15	21	1		6			132	3	135	208.03	35.33
(%)			(3.0)	(4.5)	(2.3)	(9.8)	(47.7)	(11.4)	(15.9)	(0.8)		(4.5)			(100.0)				
400～			2	15	8	9	62	20	17	4	3	8	2		150	2	152	210.27	42.41
(%)			(1.3)	(10.0)	(5.3)	(6.0)	(41.3)	(13.3)	(11.3)	(2.7)	(2.0)	(5.3)	(1.3)		(100.0)				
500～		1	3	10	4	16	74	16	25	2	2	4			157	9	166	206.75	34.46
(%)		(0.6)	(1.9)	(6.4)	(2.5)	(10.2)	(47.1)	(10.2)	(15.9)	(1.3)	(1.3)	(2.5)			(100.0)				
合計		22	153	1,048	417	1,833	6,784	2,143	2,217	234	186	621	34	15	15,707	127	15,834	209.12	35.96
(%)		(0.1)	(1.0)	(6.7)	(2.7)	(11.7)	(43.2)	(13.6)	(14.1)	(1.5)	(1.2)	(4.0)	(0.2)	(0.1)	(100.0)				
記載なし		4	18	82	19	79	155	31	32	1		5			426	12	438	187.84	34.15
(%)		(0.9)	(4.2)	(19.2)	(4.5)	(18.5)	(36.4)	(7.3)	(7.5)	(0.2)		(1.2)			(100.0)				
総計		26	171	1,130	436	1,912	6,939	2,174	2,249	235	186	626	34	15	16,133	139	16,272	208.56	36.07
(%)		(0.2)	(1.1)	(7.0)	(2.7)	(11.9)	(43.0)	(13.5)	(13.9)	(1.5)	(1.2)	(3.9)	(0.2)	(0.1)	(100.0)				
平均		142.64	129.39	106.16	109.38	94.11	104.56	107.43	107.62	116.23	109.22	112.22	101.85	88.27	105.20	166.72	105.70		
標準偏差		125.19	117.94	107.38	108.71	96.78	117.83	96.66	96.39	108.17	98.63	88.73	95.36	80.52	107.53	233.16	109.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査