

表 1975 血流量(ml/分) 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン濃度	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差
5未満		2	1	18	9	37	118	36	31	1		5	1		259	2	261	202.53	31.50
(%)		(0.8)	(0.4)	(6.9)	(3.5)	(14.3)	(45.6)	(13.9)	(12.0)	(0.4)		(1.9)	(0.4)		(100.0)				
5～		2	12	61	19	56	159	28	16		1	1			355	7	362	187.44	29.04
(%)		(0.6)	(3.4)	(17.2)	(5.4)	(15.8)	(44.8)	(7.9)	(4.5)		(0.3)	(0.3)			(100.0)				
6～	1	20	34	170	41	149	375	57	58	5	5	6		2	923	19	942	187.29	37.23
(%)	(0.1)	(2.2)	(3.7)	(18.4)	(4.4)	(16.1)	(40.6)	(6.2)	(6.3)	(0.5)	(0.5)	(0.7)		(0.2)	(100.0)				
7～	6	60	115	623	139	558	1,436	300	236	16	9	23	3	12	3,536	42	3,578	189.65	38.15
(%)	(0.2)	(1.7)	(3.3)	(17.6)	(3.9)	(15.8)	(40.6)	(8.5)	(6.7)	(0.5)	(0.3)	(0.7)	(0.1)	(0.3)	(100.0)				
8～	10	114	264	1,645	393	1,980	5,041	1,058	789	67	61	99	15	37	11,573	153	11,726	193.62	36.38
(%)	(0.1)	(1.0)	(2.3)	(14.2)	(3.4)	(17.1)	(43.6)	(9.1)	(6.8)	(0.6)	(0.5)	(0.9)	(0.1)	(0.3)	(100.0)				
9～	9	166	493	3,756	1,210	5,702	16,547	4,195	3,448	272	245	497	45	64	36,649	389	37,038	199.48	34.25
(%)	(0.0)	(0.5)	(1.3)	(10.2)	(3.3)	(15.6)	(45.1)	(11.4)	(9.4)	(0.7)	(0.7)	(1.4)	(0.1)	(0.2)	(100.0)				
10～	7	184	600	5,416	1,945	10,203	33,783	10,073	9,076	822	775	1,332	116	163	74,495	722	75,217	205.17	34.43
(%)	(0.0)	(0.2)	(0.8)	(7.3)	(2.6)	(13.7)	(45.3)	(13.5)	(12.2)	(1.1)	(1.0)	(1.8)	(0.2)	(0.2)	(100.0)				
11～	5	107	361	3,774	1,320	7,467	27,247	8,959	8,370	834	746	1,386	135	178	60,889	590	61,479	208.55	35.60
(%)	(0.0)	(0.2)	(0.6)	(6.2)	(2.2)	(12.3)	(44.7)	(14.7)	(13.7)	(1.4)	(1.2)	(2.3)	(0.2)	(0.3)	(100.0)				
12～	3	36	145	1,426	493	2,590	9,664	3,130	3,274	343	313	567	59	103	22,146	224	22,370	209.95	38.21
(%)	(0.0)	(0.2)	(0.7)	(6.4)	(2.2)	(11.7)	(43.6)	(14.1)	(14.8)	(1.5)	(1.4)	(2.6)	(0.3)	(0.5)	(100.0)				
13～		13	27	370	108	661	2,366	760	770	76	107	169	27	36	5,490	47	5,537	211.18	40.90
(%)		(0.2)	(0.5)	(6.7)	(2.0)	(12.0)	(43.1)	(13.8)	(14.0)	(1.4)	(1.9)	(3.1)	(0.5)	(0.7)	(100.0)				
14～		5	9	110	49	207	774	277	312	37	29	81	12	7	1,909	24	1,933	213.76	40.36
(%)		(0.3)	(0.5)	(5.8)	(2.6)	(10.8)	(40.5)	(14.5)	(16.3)	(1.9)	(1.5)	(4.2)	(0.6)	(0.4)	(100.0)				
合計	41	709	2,061	17,369	5,726	29,610	97,510	28,873	26,380	2,473	2,291	4,166	413	602	218,224	2,219	220,443	204.90	35.88
(%)	(0.0)	(0.3)	(0.9)	(8.0)	(2.6)	(13.6)	(44.7)	(13.2)	(12.1)	(1.1)	(1.0)	(1.9)	(0.2)	(0.3)	(100.0)				
記載なし	3	30	87	604	175	875	2,980	673	680	60	46	99	16	116	6,444	2,084	8,528	206.18	51.48
(%)	(0.0)	(0.5)	(1.4)	(9.4)	(2.7)	(13.6)	(46.2)	(10.4)	(10.6)	(0.9)	(0.7)	(1.5)	(0.2)	(1.8)	(100.0)				
総計	44	739	2,148	17,973	5,901	30,485	100,490	29,546	27,060	2,533	2,337	4,265	429	718	224,668	4,303	228,971	204.94	36.42
(%)	(0.0)	(0.3)	(1.0)	(8.0)	(2.6)	(13.6)	(44.7)	(13.2)	(12.0)	(1.1)	(1.0)	(1.9)	(0.2)	(0.3)	(100.0)				
平均	9.41	9.83	10.05	10.34	10.47	10.55	10.69	10.83	10.92	11.03	11.06	11.07	11.17	11.00	10.69	10.60	10.69		
標準偏差	1.51	1.56	1.45	1.40	1.33	1.29	1.26	1.20	1.20	1.17	1.18	1.24	1.38	1.38	1.28	1.35	1.28		

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査