

表 2007 血流量(ml/分) Non-HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

Non-HDL-C濃度	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差
50未満		2	4	32	5	35	169	86	130	16	15	22	3	4	523	6	529	222.03	41.78
(%)		(0.4)	(0.8)	(6.1)	(1.0)	(6.7)	(32.3)	(16.4)	(24.9)	(3.1)	(2.9)	(4.2)	(0.6)	(0.8)	(100.0)				
50～		2	5	76	33	151	798	416	605	54	63	151	19	8	2,381	22	2,403	225.20	39.03
(%)		(0.1)	(0.2)	(3.2)	(1.4)	(6.3)	(33.5)	(17.5)	(25.4)	(2.3)	(2.6)	(6.3)	(0.8)	(0.3)	(100.0)				
70～		2	11	149	82	378	1,720	883	1,225	161	127	304	46	26	5,114	24	5,138	224.80	39.53
(%)		(0.0)	(0.2)	(2.9)	(1.6)	(7.4)	(33.6)	(17.3)	(24.0)	(3.1)	(2.5)	(5.9)	(0.9)	(0.5)	(100.0)				
90～		7	15	222	83	422	2,083	1,138	1,450	135	191	384	58	36	6,224	41	6,265	224.81	40.71
(%)		(0.1)	(0.2)	(3.6)	(1.3)	(6.8)	(33.5)	(18.3)	(23.3)	(2.2)	(3.1)	(6.2)	(0.9)	(0.6)	(100.0)				
110～	1	4	17	160	85	367	1,713	887	1,168	138	122	363	59	54	5,138	24	5,162	226.45	44.09
(%)	(0.0)	(0.1)	(0.3)	(3.1)	(1.7)	(7.1)	(33.3)	(17.3)	(22.7)	(2.7)	(2.4)	(7.1)	(1.1)	(1.1)	(100.0)				
130～			9	104	41	242	1,015	560	691	106	96	211	31	26	3,132	17	3,149	226.12	41.93
(%)			(0.3)	(3.3)	(1.3)	(7.7)	(32.4)	(17.9)	(22.1)	(3.4)	(3.1)	(6.7)	(1.0)	(0.8)	(100.0)				
150～		1	2	35	24	118	573	262	330	44	62	114	10	15	1,590	7	1,597	226.54	42.74
(%)		(0.1)	(0.1)	(2.2)	(1.5)	(7.4)	(36.0)	(16.5)	(20.8)	(2.8)	(3.9)	(7.2)	(0.6)	(0.9)	(100.0)				
170～		1	1	28	10	57	238	110	135	18	28	40	8	3	677	5	682	223.18	41.75
(%)		(0.1)	(0.1)	(4.1)	(1.5)	(8.4)	(35.2)	(16.2)	(19.9)	(2.7)	(4.1)	(5.9)	(1.2)	(0.4)	(100.0)				
190～		1		11	6	25	86	50	50	4	8	13	4	2	260	2	262	222.00	44.62
(%)		(0.4)		(4.2)	(2.3)	(9.6)	(33.1)	(19.2)	(19.2)	(1.5)	(3.1)	(5.0)	(1.5)	(0.8)	(100.0)				
210～		1		3	2	15	46	25	32	7	1	5	1		138	3	141	220.07	36.19
(%)		(0.7)		(2.2)	(1.4)	(10.9)	(33.3)	(18.1)	(23.2)	(5.1)	(0.7)	(3.6)	(0.7)		(100.0)				
合計	1	21	64	820	371	1,810	8,441	4,417	5,816	683	713	1,607	239	174	25,177	151	25,328	225.30	41.39
(%)	(0.0)	(0.1)	(0.3)	(3.3)	(1.5)	(7.2)	(33.5)	(17.5)	(23.1)	(2.7)	(2.8)	(6.4)	(0.9)	(0.7)	(100.0)				
記載なし		16	30	522	207	1,136	5,106	2,190	3,311	395	307	858	106	28	14,212	406	14,618	222.20	38.37
(%)		(0.1)	(0.2)	(3.7)	(1.5)	(8.0)	(35.9)	(15.4)	(23.3)	(2.8)	(2.2)	(6.0)	(0.7)	(0.2)	(100.0)				
総計	1	37	94	1,342	578	2,946	13,547	6,607	9,127	1,078	1,020	2,465	345	202	39,389	557	39,946	224.18	40.35
(%)	(0.0)	(0.1)	(0.2)	(3.4)	(1.5)	(7.5)	(34.4)	(16.8)	(23.2)	(2.7)	(2.6)	(6.3)	(0.9)	(0.5)	(100.0)				
平均	117.00	109.33	104.50	105.95	108.23	109.33	107.25	107.02	105.38	108.11	110.13	108.29	109.40	113.33	107.13	105.79	107.12		
標準偏差		44.30	30.77	33.74	33.99	34.52	33.60	33.08	32.83	34.72	34.28	31.76	32.09	29.97	33.33	39.56	33.37		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査