

表 2017 HDF希釈方法 原疾患別（オフラインHDF 透析回数週3回）

原疾患	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
慢性糸球体腎炎	287	1,782	2	3	2,074		376	2,450
(%)	(13.8)	(85.9)	(0.1)	(0.1)	(100.0)			
慢性腎盂腎炎	3	53		1	57		7	64
(%)	(5.3)	(93.0)		(1.8)	(100.0)			
急速進行性糸球体腎炎	3	29			32		8	40
(%)	(9.4)	(90.6)			(100.0)			
妊娠腎/妊娠中毒症	4	51			55		11	66
(%)	(7.3)	(92.7)			(100.0)			
その他の分類不能の腎炎	6	19			25		5	30
(%)	(24.0)	(76.0)			(100.0)			
多発性嚢胞腎	19	125			144		44	188
(%)	(13.2)	(86.8)			(100.0)			
腎硬化症	31	233	1		265		71	336
(%)	(11.7)	(87.9)	(0.4)		(100.0)			
悪性高血圧	5	30			35		9	44
(%)	(14.3)	(85.7)			(100.0)			
糖尿病性腎症	191	1,260	1	1	1,453		290	1,743
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
SLE腎炎	7	30			37		15	52
(%)	(18.9)	(81.1)			(100.0)			
アミロイド腎		7			7		1	8
(%)		(100.0)			(100.0)			
痛風腎	1	18			19		6	25
(%)	(5.3)	(94.7)			(100.0)			
先天性代謝異常による腎不全	1	4			5		1	6
(%)	(20.0)	(80.0)			(100.0)			
腎・尿路結核	1	2			3		1	4
(%)	(33.3)	(66.7)			(100.0)			
腎・尿路結石	1	10			11		1	12
(%)	(9.1)	(90.9)			(100.0)			
腎・尿路腫瘍	1	12			13		4	17
(%)	(7.7)	(92.3)			(100.0)			
閉塞性尿路障害	4	4			8		2	10
(%)	(50.0)	(50.0)			(100.0)			
骨髄腫	1	6			7		1	8
(%)	(14.3)	(85.7)			(100.0)			
腎形成不全	1	10			11		2	13
(%)	(9.1)	(90.9)			(100.0)			
不明	43	358	1		402		110	512
(%)	(10.7)	(89.1)	(0.2)		(100.0)			
再導入	9	54			63		9	72
(%)	(14.3)	(85.7)			(100.0)			
その他	15	97			112		16	128
(%)	(13.4)	(86.6)			(100.0)			
合計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
記載なし								
(%)								
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査