

表 2029 HDF希釈方法 身長(cm)別（オフラインHDF 透析回数週3回）

身長	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
130未満	2	5			7		3	10
(%)	(28.6)	(71.4)			(100.0)			
130～		6			6		4	10
(%)		(100.0)			(100.0)			
135～	2	40			42		9	51
(%)	(4.8)	(95.2)			(100.0)			
140～	27	160			187		30	217
(%)	(14.4)	(85.6)			(100.0)			
145～	55	365			420		92	512
(%)	(13.1)	(86.9)			(100.0)			
150～	96	622			718		132	850
(%)	(13.4)	(86.6)			(100.0)			
155～	101	666	2	2	771		158	929
(%)	(13.1)	(86.4)	(0.3)	(0.3)	(100.0)			
160～	112	718	2	1	833		142	975
(%)	(13.4)	(86.2)	(0.2)	(0.1)	(100.0)			
165～	105	680		1	786		142	928
(%)	(13.4)	(86.5)		(0.1)	(100.0)			
170～	79	471	1	1	552		108	660
(%)	(14.3)	(85.3)	(0.2)	(0.2)	(100.0)			
175～	19	187			206		45	251
(%)	(9.2)	(90.8)			(100.0)			
180～	9	51			60		7	67
(%)	(15.0)	(85.0)			(100.0)			
185～	1	10			11		4	15
(%)	(9.1)	(90.9)			(100.0)			
190～		1			1		1	2
(%)		(100.0)			(100.0)			
合計	608	3,982	5	5	4,600		877	5,477
(%)	(13.2)	(86.6)	(0.1)	(0.1)	(100.0)			
記載なし	26	212			238		113	351
(%)	(10.9)	(89.1)			(100.0)			
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
平均	159.91	159.96	162.00	163.60	159.96		159.74	159.92
標準偏差	9.54	9.72	6.89	5.55	9.69		10.33	9.79

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査