

表 2030 HDF希釈方法 透析後体重(kg)別（オフラインHDF 透析回数週3回）

透析後体重	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
20未満							1	1
(%)								
20～	5	11			16		3	19
(%)	(31.3)	(68.8)			(100.0)			
30～	51	389	1		441		101	542
(%)	(11.6)	(88.2)	(0.2)		(100.0)			
40～	201	1,172	1	1	1,375		271	1,646
(%)	(14.6)	(85.2)	(0.1)	(0.1)	(100.0)			
50～	200	1,311	2	3	1,516		300	1,816
(%)	(13.2)	(86.5)	(0.1)	(0.2)	(100.0)			
60～	104	787		1	892		186	1,078
(%)	(11.7)	(88.2)		(0.1)	(100.0)			
70～	47	276			323		51	374
(%)	(14.6)	(85.4)			(100.0)			
80～	12	105			117		29	146
(%)	(10.3)	(89.7)			(100.0)			
90～	2	45	1		48		5	53
(%)	(4.2)	(93.8)	(2.1)		(100.0)			
100～	2	11			13			13
(%)	(15.4)	(84.6)			(100.0)			
110～		8			8		1	9
(%)		(100.0)			(100.0)			
120～	1	6			7			7
(%)	(14.3)	(85.7)			(100.0)			
合計	625	4,121	5	5	4,756		948	5,704
(%)	(13.1)	(86.6)	(0.1)	(0.1)	(100.0)			
記載なし	9	73			82		42	124
(%)	(11.0)	(89.0)			(100.0)			
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
平均	54.14	54.94	58.78	52.78	54.84		54.08	54.71
標準偏差	12.02	13.02	23.08	7.97	12.90		12.01	12.76

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査