

表 2033 HDF希釈方法 Body Mass Index別（オフラインHDF 透析回数週3回）

Body Mass Index	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
14未満	4	21			25		10	35
(%)	(16.0)	(84.0)			(100.0)			
14～	27	161	1		189		30	219
(%)	(14.3)	(85.2)	(0.5)		(100.0)			
16～	80	525	1	1	607		118	725
(%)	(13.2)	(86.5)	(0.2)	(0.2)	(100.0)			
18～	152	910		1	1,063		199	1,262
(%)	(14.3)	(85.6)		(0.1)	(100.0)			
20～	127	873	1	2	1,003		192	1,195
(%)	(12.7)	(87.0)	(0.1)	(0.2)	(100.0)			
22～	91	633	1	1	726		162	888
(%)	(12.5)	(87.2)	(0.1)	(0.1)	(100.0)			
24～	55	394			449		76	525
(%)	(12.2)	(87.8)			(100.0)			
26～	36	197			233		34	267
(%)	(15.5)	(84.5)			(100.0)			
28～	16	106			122		20	142
(%)	(13.1)	(86.9)			(100.0)			
30～	11	91	1		103		13	116
(%)	(10.7)	(88.3)	(1.0)		(100.0)			
合計	599	3,911	5	5	4,520		854	5,374
(%)	(13.3)	(86.5)	(0.1)	(0.1)	(100.0)			
記載なし	35	283			318		136	454
(%)	(11.0)	(89.0)			(100.0)			
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
平均	21.07	21.19	22.01	19.65	21.17		20.95	21.14
標準偏差	3.62	3.60	6.68	2.13	3.61		3.42	3.58

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査