

表 2034 HDF希釈方法 透析前BUN(mg/dL)別（オフラインHDF 透析回数週3回 透析歴2年以上）

透析前BUN	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
20未満	1	8			9		2	11
(%)	(11.1)	(88.9)			(100.0)			
20～	12	41			53		11	64
(%)	(22.6)	(77.4)			(100.0)			
30～	22	188			210		41	251
(%)	(10.5)	(89.5)			(100.0)			
40～	68	543			611		127	738
(%)	(11.1)	(88.9)			(100.0)			
50～	145	946	1	3	1,095		193	1,288
(%)	(13.2)	(86.4)	(0.1)	(0.3)	(100.0)			
60～	152	973	2		1,127		182	1,309
(%)	(13.5)	(86.3)	(0.2)		(100.0)			
70～	91	664		1	756		178	934
(%)	(12.0)	(87.8)		(0.1)	(100.0)			
80～	39	292			331		74	405
(%)	(11.8)	(88.2)			(100.0)			
90～	23	119	2		144		15	159
(%)	(16.0)	(82.6)	(1.4)		(100.0)			
100～	5	36			41		4	45
(%)	(12.2)	(87.8)			(100.0)			
110～	1	3			4		2	6
(%)	(25.0)	(75.0)			(100.0)			
120～		4			4			4
(%)		(100.0)			(100.0)			
合計	559	3,817	5	4	4,385		829	5,214
(%)	(12.7)	(87.0)	(0.1)	(0.1)	(100.0)			
記載なし	6	62			68		38	106
(%)	(8.8)	(91.2)			(100.0)			
総計	565	3,879	5	4	4,453		867	5,320
(%)	(12.7)	(87.1)	(0.1)	(0.1)	(100.0)			
平均	62.37	61.83	74.00	60.00	61.91		61.71	61.88
標準偏差	15.57	15.35	19.14	8.16	15.38		15.25	15.36

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査