

表 2041 HDF希釈方法 透析前CRP濃度(mg/dL)別（オフラインHDF 透析回数週3回）

透析前CRP濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
0.2未満	277	1,936	2	3	2,218		424	2,642
(%)	(12.5)	(87.3)	(0.1)	(0.1)	(100.0)			
0.2～	92	692		1	785		148	933
(%)	(11.7)	(88.2)		(0.1)	(100.0)			
0.5～	60	356			416		76	492
(%)	(14.4)	(85.6)			(100.0)			
1.0～	33	274			307		50	357
(%)	(10.7)	(89.3)			(100.0)			
2.0～	25	171			196		45	241
(%)	(12.8)	(87.2)			(100.0)			
4.0～	13	57			70		18	88
(%)	(18.6)	(81.4)			(100.0)			
6.0～	5	22			27		8	35
(%)	(18.5)	(81.5)			(100.0)			
8.0～	2	16			18			18
(%)	(11.1)	(88.9)			(100.0)			
10.0～	4	18			22		4	26
(%)	(18.2)	(81.8)			(100.0)			
15.0～	2	19			21		1	22
(%)	(9.5)	(90.5)			(100.0)			
合計	513	3,561	2	4	4,080		774	4,854
(%)	(12.6)	(87.3)	(0.0)	(0.1)	(100.0)			
記載なし	121	633	3	1	758		216	974
(%)	(16.0)	(83.5)	(0.4)	(0.1)	(100.0)			
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
平均	0.83	0.74	0.12	0.14	0.75		0.71	0.75
標準偏差	2.46	2.09	0.03	0.19	2.14		1.93	2.10

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査