

表 2042 HDF希釈方法 透析前補正カルシウム濃度(mg/dL)別（オフラインHDF 透析回数週3回）

透析前補正 カルシウム濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
7.0未満	1	25			26		1	27
(%)	(3.8)	(96.2)			(100.0)			
7.0～	2	26			28		5	33
(%)	(7.1)	(92.9)			(100.0)			
7.5～	17	81			98		25	123
(%)	(17.3)	(82.7)			(100.0)			
8.0～	38	274	1		313		69	382
(%)	(12.1)	(87.5)	(0.3)		(100.0)			
8.4～	12	85			97		15	112
(%)	(12.4)	(87.6)			(100.0)			
8.5～	128	808		1	937		232	1,169
(%)	(13.7)	(86.2)		(0.1)	(100.0)			
9.0～	169	1,184		1	1,354		231	1,585
(%)	(12.5)	(87.4)		(0.1)	(100.0)			
9.5～	138	858		3	999		176	1,175
(%)	(13.8)	(85.9)		(0.3)	(100.0)			
10.0～	17	101	1		119		25	144
(%)	(14.3)	(84.9)	(0.8)		(100.0)			
10.1～	47	342			389		90	479
(%)	(12.1)	(87.9)			(100.0)			
10.5～	25	182			207		32	239
(%)	(12.1)	(87.9)			(100.0)			
11.0～	12	59			71		10	81
(%)	(16.9)	(83.1)			(100.0)			
11.5～	8	22			30		3	33
(%)	(26.7)	(73.3)			(100.0)			
12.0～	2	17			19		3	22
(%)	(10.5)	(89.5)			(100.0)			
合計	616	4,064	2	5	4,687		917	5,604
(%)	(13.1)	(86.7)	(0.0)	(0.1)	(100.0)			
記載なし	18	130	3		151		73	224
(%)	(11.9)	(86.1)	(2.0)		(100.0)			
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
平均	9.34	9.29	9.05	9.36	9.29		9.25	9.28
標準偏差	0.80	0.82	1.34	0.52	0.82		0.78	0.81

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査