

表 2043 HDF希釈方法 透析前リン濃度(mg/dL)別（オフラインHDF 透析回数週3回）

透析前リン濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
3未満	28	178			206		35	241
(%)	(13.6)	(86.4)			(100.0)			
3.0～	26	188			214		46	260
(%)	(12.1)	(87.9)			(100.0)			
3.5～	46	327			373		78	451
(%)	(12.3)	(87.7)			(100.0)			
4.0～	75	491	1	2	569		112	681
(%)	(13.2)	(86.3)	(0.2)	(0.4)	(100.0)			
4.5～	90	578	1	1	670		151	821
(%)	(13.4)	(86.3)	(0.1)	(0.1)	(100.0)			
5.0～	97	663	1	1	762		154	916
(%)	(12.7)	(87.0)	(0.1)	(0.1)	(100.0)			
5.5～	86	538		1	625		108	733
(%)	(13.8)	(86.1)		(0.2)	(100.0)			
6.0～	11	80			91		19	110
(%)	(12.1)	(87.9)			(100.0)			
6.1～	51	307			358		65	423
(%)	(14.2)	(85.8)			(100.0)			
6.5～	40	271			311		60	371
(%)	(12.9)	(87.1)			(100.0)			
7.0～	27	178			205		38	243
(%)	(13.2)	(86.8)			(100.0)			
7.5～	16	116			132		30	162
(%)	(12.1)	(87.9)			(100.0)			
8.0～	11	68			79		9	88
(%)	(13.9)	(86.1)			(100.0)			
8.5～	5	53			58		9	67
(%)	(8.6)	(91.4)			(100.0)			
9.0～	14	76	2		92		17	109
(%)	(15.2)	(82.6)	(2.2)		(100.0)			
合計	623	4,112	5	5	4,745		931	5,676
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
記載なし	11	82			93		59	152
(%)	(11.8)	(88.2)			(100.0)			
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
平均	5.29	5.28	6.66	4.82	5.28		5.24	5.27
標準偏差	1.51	1.51	2.73	0.76	1.51		1.44	1.50

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査