

表 2046 HDF希釈方法 透析前ヘモグロビン濃度(g/dL)別（オフラインHDF 透析回数週3回）

透析前ヘモグロビン 濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
5未満	1	7			8		2	10
(%)	(12.5)	(87.5)			(100.0)			
5～	2	5			7		2	9
(%)	(28.6)	(71.4)			(100.0)			
6～	6	32			38		4	42
(%)	(15.8)	(84.2)			(100.0)			
7～	11	78			89		17	106
(%)	(12.4)	(87.6)			(100.0)			
8～	23	267		1	291		58	349
(%)	(7.9)	(91.8)		(0.3)	(100.0)			
9～	110	690	2	1	803		156	959
(%)	(13.7)	(85.9)	(0.2)	(0.1)	(100.0)			
10～	205	1,282	1	1	1,489		296	1,785
(%)	(13.8)	(86.1)	(0.1)	(0.1)	(100.0)			
11～	169	1,120		2	1,291		258	1,549
(%)	(13.1)	(86.8)		(0.2)	(100.0)			
12～	62	473	2		537		74	611
(%)	(11.5)	(88.1)	(0.4)		(100.0)			
13～	18	104			122		20	142
(%)	(14.8)	(85.2)			(100.0)			
14～	20	45			65		17	82
(%)	(30.8)	(69.2)			(100.0)			
合計	627	4,103	5	5	4,740		904	5,644
(%)	(13.2)	(86.6)	(0.1)	(0.1)	(100.0)			
記載なし	7	91			98		86	184
(%)	(7.1)	(92.9)			(100.0)			
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
平均	10.78	10.68	10.62	10.34	10.69		10.66	10.68
標準偏差	1.51	1.36	1.49	1.01	1.38		1.43	1.39

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査