

表 2049 HDF希釈方法 Non-HDL-C濃度(mg/dL)別（オフラインHDF 透析回数週3回）

Non-HDL-C濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
50未満	10	55			65		11	76
(%)	(15.4)	(84.6)			(100.0)			
50～	39	252			291		57	348
(%)	(13.4)	(86.6)			(100.0)			
70～	94	488	1	1	584		127	711
(%)	(16.1)	(83.6)	(0.2)	(0.2)	(100.0)			
90～	120	592	1		713		135	848
(%)	(16.8)	(83.0)	(0.1)		(100.0)			
110～	91	455			546		118	664
(%)	(16.7)	(83.3)			(100.0)			
130～	40	304	1		345		58	403
(%)	(11.6)	(88.1)	(0.3)		(100.0)			
150～	34	129			163		30	193
(%)	(20.9)	(79.1)			(100.0)			
170～	8	62		1	71		14	85
(%)	(11.3)	(87.3)		(1.4)	(100.0)			
190～	3	15			18		4	22
(%)	(16.7)	(83.3)			(100.0)			
210～	3	17	1		21		5	26
(%)	(14.3)	(81.0)	(4.8)		(100.0)			
合計	442	2,369	4	2	2,817		559	3,376
(%)	(15.7)	(84.1)	(0.1)	(0.1)	(100.0)			
記載なし	192	1,825	1	3	2,021		431	2,452
(%)	(9.5)	(90.3)	(0.0)	(0.1)	(100.0)			
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
平均	105.62	105.57	134.25	134.50	105.64		105.10	105.55
標準偏差	32.32	33.11	54.65	71.42	33.05		33.63	33.14

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査