

表 2052 HDF希釈方法 透析前脈拍(拍/分)別 (オフラインHDF 透析回数週3回)

透析前脈拍	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
40未満	1	3			4			4
(%)	(25.0)	(75.0)			(100.0)			
40～	6	17			23		10	33
(%)	(26.1)	(73.9)			(100.0)			
50～	42	248			290		50	340
(%)	(14.5)	(85.5)			(100.0)			
60～	129	864	2	2	997		158	1,155
(%)	(12.9)	(86.7)	(0.2)	(0.2)	(100.0)			
70～	186	1,087	3	2	1,278		213	1,491
(%)	(14.6)	(85.1)	(0.2)	(0.2)	(100.0)			
80～	111	672		1	784		185	969
(%)	(14.2)	(85.7)		(0.1)	(100.0)			
90～	49	342			391		75	466
(%)	(12.5)	(87.5)			(100.0)			
100～	9	130			139		14	153
(%)	(6.5)	(93.5)			(100.0)			
110～	4	37			41		7	48
(%)	(9.8)	(90.2)			(100.0)			
120～		15			15		5	20
(%)		(100.0)			(100.0)			
合計	537	3,415	5	5	3,962		717	4,679
(%)	(13.6)	(86.2)	(0.1)	(0.1)	(100.0)			
記載なし	97	779			876		273	1,149
(%)	(11.1)	(88.9)			(100.0)			
総計	634	4,194	5	5	4,838		990	5,828
(%)	(13.1)	(86.7)	(0.1)	(0.1)	(100.0)			
平均	74.58	75.79	68.80	71.00	75.61		76.32	75.72
標準偏差	11.93	13.39	7.56	7.42	13.20		13.20	13.20

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査