

表 2074 HDF希釈方法 原疾患別（オンラインHDF 透析回数週3回）

原疾患	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
慢性糸球体腎炎	10,615	540		396	11,551		813	12,364
(%)	(91.9)	(4.7)		(3.4)	(100.0)			
慢性腎盂腎炎	300	16		14	330		19	349
(%)	(90.9)	(4.8)		(4.2)	(100.0)			
急速進行性糸球体腎炎	211	4		7	222		14	236
(%)	(95.0)	(1.8)		(3.2)	(100.0)			
妊娠腎/妊娠中毒症	249	12		9	270		18	288
(%)	(92.2)	(4.4)		(3.3)	(100.0)			
その他の分類不能の腎炎	141	7		3	151		13	164
(%)	(93.4)	(4.6)		(2.0)	(100.0)			
多発性嚢胞腎	1,092	47		31	1,170		82	1,252
(%)	(93.3)	(4.0)		(2.6)	(100.0)			
腎硬化症	2,140	119		100	2,359		122	2,481
(%)	(90.7)	(5.0)		(4.2)	(100.0)			
悪性高血圧	232	9		9	250		15	265
(%)	(92.8)	(3.6)		(3.6)	(100.0)			
糖尿病性腎症	9,651	408		522	10,581	1	518	11,100
(%)	(91.2)	(3.9)		(4.9)	(100.0)			
SLE腎炎	263	14		7	284		18	302
(%)	(92.6)	(4.9)		(2.5)	(100.0)			
アミロイド腎	42	6			48		2	50
(%)	(87.5)	(12.5)			(100.0)			
痛風腎	106	8		3	117		10	127
(%)	(90.6)	(6.8)		(2.6)	(100.0)			
先天性代謝異常による腎不全	32	3		2	37		1	38
(%)	(86.5)	(8.1)		(5.4)	(100.0)			
腎・尿路結核	14	1		1	16		2	18
(%)	(87.5)	(6.3)		(6.3)	(100.0)			
腎・尿路結石	49	1		2	52		3	55
(%)	(94.2)	(1.9)		(3.8)	(100.0)			
腎・尿路腫瘍	52	3		6	61		4	65
(%)	(85.2)	(4.9)		(9.8)	(100.0)			
閉塞性尿路障害	57	2		1	60		2	62
(%)	(95.0)	(3.3)		(1.7)	(100.0)			
骨髄腫	11				11			11
(%)	(100.0)				(100.0)			
腎形成不全	73	3		5	81		5	86
(%)	(90.1)	(3.7)		(6.2)	(100.0)			
不明	2,557	83		131	2,771		153	2,924
(%)	(92.3)	(3.0)		(4.7)	(100.0)			
再導入	293	18		11	322		17	339
(%)	(91.0)	(5.6)		(3.4)	(100.0)			
その他	603	22		20	645		79	724
(%)	(93.5)	(3.4)		(3.1)	(100.0)			
合計	28,783	1,326		1,280	31,389	1	1,910	33,300
(%)	(91.7)	(4.2)		(4.1)	(100.0)			
記載なし								
(%)								
総計	28,783	1,326		1,280	31,389	1	1,910	33,300
(%)	(91.7)	(4.2)		(4.1)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査