

表 2086 HDF希釈方法 身長(cm)別 (オンラインHDF 透析回数週3回)

身長	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
130未満	27				27		1	28
(%)	(100.0)				(100.0)			
130～	51			2	53		5	58
(%)	(96.2)			(3.8)	(100.0)			
135～	194	10		13	217		13	230
(%)	(89.4)	(4.6)		(6.0)	(100.0)			
140～	785	36		48	869		41	910
(%)	(90.3)	(4.1)		(5.5)	(100.0)			
145～	2,064	108		106	2,278		133	2,411
(%)	(90.6)	(4.7)		(4.7)	(100.0)			
150～	3,666	183		169	4,018		215	4,233
(%)	(91.2)	(4.6)		(4.2)	(100.0)			
155～	4,374	201		196	4,771		275	5,046
(%)	(91.7)	(4.2)		(4.1)	(100.0)			
160～	5,262	258		233	5,753		302	6,055
(%)	(91.5)	(4.5)		(4.1)	(100.0)			
165～	4,957	229		196	5,382	1	280	5,663
(%)	(92.1)	(4.3)		(3.6)	(100.0)			
170～	3,551	170		111	3,832		202	4,034
(%)	(92.7)	(4.4)		(2.9)	(100.0)			
175～	1,336	63		29	1,428		79	1,507
(%)	(93.6)	(4.4)		(2.0)	(100.0)			
180～	373	15		13	401		18	419
(%)	(93.0)	(3.7)		(3.2)	(100.0)			
185～	54	2			56		6	62
(%)	(96.4)	(3.6)			(100.0)			
190～	4	1		2	7			7
(%)	(57.1)	(14.3)		(28.6)	(100.0)			
合計	26,698	1,276		1,118	29,092	1	1,570	30,663
(%)	(91.8)	(4.4)		(3.8)	(100.0)			
記載なし	2,085	50		162	2,297		340	2,637
(%)	(90.8)	(2.2)		(7.1)	(100.0)			
総計	28,783	1,326		1,280	31,389	1	1,910	33,300
(%)	(91.7)	(4.2)		(4.1)	(100.0)			
平均	160.99	160.85		159.39	160.92	168.00	160.83	160.91
標準偏差	9.49	9.25		9.45	9.48		9.38	9.48

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査