

表 2087 HDF希釈方法 透析後体重(kg)別（オンラインHDF 透析回数週3回）

透析後体重	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
20未満	8	2		1	11			11
(%)	(72.7)	(18.2)		(9.1)	(100.0)			
20～	45	1		5	51		3	54
(%)	(88.2)	(2.0)		(9.8)	(100.0)			
30～	1,747	86		89	1,922		131	2,053
(%)	(90.9)	(4.5)		(4.6)	(100.0)			
40～	6,986	336		335	7,657		499	8,156
(%)	(91.2)	(4.4)		(4.4)	(100.0)			
50～	9,055	417		391	9,863	1	555	10,419
(%)	(91.8)	(4.2)		(4.0)	(100.0)			
60～	6,233	247		234	6,714		407	7,121
(%)	(92.8)	(3.7)		(3.5)	(100.0)			
70～	2,624	113		81	2,818		170	2,988
(%)	(93.1)	(4.0)		(2.9)	(100.0)			
80～	988	49		19	1,056		47	1,103
(%)	(93.6)	(4.6)		(1.8)	(100.0)			
90～	353	17		7	377		21	398
(%)	(93.6)	(4.5)		(1.9)	(100.0)			
100～	122	1		2	125		4	129
(%)	(97.6)	(0.8)		(1.6)	(100.0)			
110～	47			3	50		2	52
(%)	(94.0)			(6.0)	(100.0)			
120～	27	1		1	29		1	30
(%)	(93.1)	(3.4)		(3.4)	(100.0)			
合計	28,235	1,270		1,168	30,673	1	1,840	32,514
(%)	(92.1)	(4.1)		(3.8)	(100.0)			
記載なし	548	56		112	716		70	786
(%)	(76.5)	(7.8)		(15.6)	(100.0)			
総計	28,783	1,326		1,280	31,389	1	1,910	33,300
(%)	(91.7)	(4.2)		(4.1)	(100.0)			
平均	57.18	56.37		54.82	57.06	51.80	56.22	57.01
標準偏差	13.13	12.92		12.33	13.10		12.53	13.07

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査