

表 2090 HDF希釈方法 Body Mass Index別（オンラインHDF 透析回数週3回）

Body Mass Index	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
14未満	88	3		3	94		11	105
(%)	(93.6)	(3.2)		(3.2)	(100.0)			
14～	682	33		46	761		41	802
(%)	(89.6)	(4.3)		(6.0)	(100.0)			
16～	2,743	143		112	2,998		176	3,174
(%)	(91.5)	(4.8)		(3.7)	(100.0)			
18～	5,370	280		230	5,880	1	324	6,205
(%)	(91.3)	(4.8)		(3.9)	(100.0)			
20～	6,168	274		277	6,719		376	7,095
(%)	(91.8)	(4.1)		(4.1)	(100.0)			
22～	4,835	231		198	5,264		278	5,542
(%)	(91.9)	(4.4)		(3.8)	(100.0)			
24～	3,061	109		145	3,315		170	3,485
(%)	(92.3)	(3.3)		(4.4)	(100.0)			
26～	1,687	66		56	1,809		84	1,893
(%)	(93.3)	(3.6)		(3.1)	(100.0)			
28～	890	48		22	960		47	1,007
(%)	(92.7)	(5.0)		(2.3)	(100.0)			
30～	805	32		19	856		35	891
(%)	(94.0)	(3.7)		(2.2)	(100.0)			
合計	26,329	1,219		1,108	28,656	1	1,542	30,199
(%)	(91.9)	(4.3)		(3.9)	(100.0)			
記載なし	2,454	107		172	2,733		368	3,101
(%)	(89.8)	(3.9)		(6.3)	(100.0)			
総計	28,783	1,326		1,280	31,389	1	1,910	33,300
(%)	(91.7)	(4.2)		(4.1)	(100.0)			
平均	21.81	21.53		21.45	21.79	18.35	21.51	21.77
標準偏差	3.67	3.63		3.41	3.66		3.55	3.65

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査