

表 2091 HDF希釈方法 透析前BUN(mg/dL)別（オンラインHDF 透析回数週3回 透析歴2年以上）

透析前BUN	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
20未満	20			3	23		4	27
(%)	(87.0)			(13.0)	(100.0)			
20～	204	7		7	218		13	231
(%)	(93.6)	(3.2)		(3.2)	(100.0)			
30～	1,177	46		41	1,264		83	1,347
(%)	(93.1)	(3.6)		(3.2)	(100.0)			
40～	3,426	155		108	3,689		222	3,911
(%)	(92.9)	(4.2)		(2.9)	(100.0)			
50～	5,976	280		228	6,484	1	375	6,860
(%)	(92.2)	(4.3)		(3.5)	(100.0)			
60～	6,369	322		222	6,913		431	7,344
(%)	(92.1)	(4.7)		(3.2)	(100.0)			
70～	4,419	197		154	4,770		282	5,052
(%)	(92.6)	(4.1)		(3.2)	(100.0)			
80～	2,075	96		91	2,262		126	2,388
(%)	(91.7)	(4.2)		(4.0)	(100.0)			
90～	746	34		37	817		63	880
(%)	(91.3)	(4.2)		(4.5)	(100.0)			
100～	205	7		8	220		8	228
(%)	(93.2)	(3.2)		(3.6)	(100.0)			
110～	58	2		2	62		5	67
(%)	(93.5)	(3.2)		(3.2)	(100.0)			
120～	12	1		2	15		1	16
(%)	(80.0)	(6.7)		(13.3)	(100.0)			
合計	24,687	1,147		903	26,737	1	1,613	28,351
(%)	(92.3)	(4.3)		(3.4)	(100.0)			
記載なし	375	43		98	516		89	605
(%)	(72.7)	(8.3)		(19.0)	(100.0)			
総計	25,062	1,190		1,001	27,253	1	1,702	28,956
(%)	(92.0)	(4.4)		(3.7)	(100.0)			
平均	62.36	62.52		63.35	62.40	58.00	62.32	62.40
標準偏差	14.99	14.49		15.88	15.00		15.22	15.01

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査