

表 2099 HDF希釈方法 透析前補正カルシウム濃度(mg/dL)別（オンラインHDF 透析回数週3回）

透析前補正 カルシウム濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
7.0未満	112	4		7	123		7	130
(%)	(91.1)	(3.3)		(5.7)	(100.0)			
7.0～	197	4		13	214		7	221
(%)	(92.1)	(1.9)		(6.1)	(100.0)			
7.5～	704	33		27	764		53	817
(%)	(92.1)	(4.3)		(3.5)	(100.0)			
8.0～	2,111	85		111	2,307	1	128	2,436
(%)	(91.5)	(3.7)		(4.8)	(100.0)			
8.4～	606	25		27	658		35	693
(%)	(92.1)	(3.8)		(4.1)	(100.0)			
8.5～	6,955	257		320	7,532		413	7,945
(%)	(92.3)	(3.4)		(4.2)	(100.0)			
9.0～	8,532	394		330	9,256		458	9,714
(%)	(92.2)	(4.3)		(3.6)	(100.0)			
9.5～	5,453	261		198	5,912		351	6,263
(%)	(92.2)	(4.4)		(3.3)	(100.0)			
10.0～	631	34		19	684		53	737
(%)	(92.3)	(5.0)		(2.8)	(100.0)			
10.1～	1,677	99		59	1,835		95	1,930
(%)	(91.4)	(5.4)		(3.2)	(100.0)			
10.5～	803	43		31	877		32	909
(%)	(91.6)	(4.9)		(3.5)	(100.0)			
11.0～	257	13		12	282		18	300
(%)	(91.1)	(4.6)		(4.3)	(100.0)			
11.5～	87	2		3	92		4	96
(%)	(94.6)	(2.2)		(3.3)	(100.0)			
12.0～	36	3		6	45		1	46
(%)	(80.0)	(6.7)		(13.3)	(100.0)			
合計	28,161	1,257		1,163	30,581	1	1,655	32,237
(%)	(92.1)	(4.1)		(3.8)	(100.0)			
記載なし	622	69		117	808		255	1,063
(%)	(77.0)	(8.5)		(14.5)	(100.0)			
総計	28,783	1,326		1,280	31,389	1	1,910	33,300
(%)	(91.7)	(4.2)		(4.1)	(100.0)			
平均	9.17	9.25		9.11	9.17	8.30	9.16	9.17
標準偏差	0.73	0.75		0.80	0.73		0.72	0.73

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査