

表 2103 HDF希釈方法 透析前ヘモグロビン濃度(g/dL)別（オンラインHDF 透析回数週3回）

透析前ヘモグロビン濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
5未満	14	1			15		1	16
(%)	(93.3)	(6.7)			(100.0)			
5～	18	1		1	20		1	21
(%)	(90.0)	(5.0)		(5.0)	(100.0)			
6～	57	7		3	67		2	69
(%)	(85.1)	(10.4)		(4.5)	(100.0)			
7～	282	15		13	310		18	328
(%)	(91.0)	(4.8)		(4.2)	(100.0)			
8～	1,059	57		52	1,168		76	1,244
(%)	(90.7)	(4.9)		(4.5)	(100.0)			
9～	4,128	186		220	4,534		242	4,776
(%)	(91.0)	(4.1)		(4.9)	(100.0)			
10～	9,459	419		423	10,301	1	593	10,895
(%)	(91.8)	(4.1)		(4.1)	(100.0)			
11～	8,587	395		284	9,266		516	9,782
(%)	(92.7)	(4.3)		(3.1)	(100.0)			
12～	3,255	133		114	3,502		181	3,683
(%)	(92.9)	(3.8)		(3.3)	(100.0)			
13～	883	39		40	962		63	1,025
(%)	(91.8)	(4.1)		(4.2)	(100.0)			
14～	390	9		12	411		17	428
(%)	(94.9)	(2.2)		(2.9)	(100.0)			
合計	28,132	1,262		1,162	30,556	1	1,710	32,267
(%)	(92.1)	(4.1)		(3.8)	(100.0)			
記載なし	651	64		118	833		200	1,033
(%)	(78.2)	(7.7)		(14.2)	(100.0)			
総計	28,783	1,326		1,280	31,389	1	1,910	33,300
(%)	(91.7)	(4.2)		(4.1)	(100.0)			
平均	10.88	10.80		10.71	10.87	10.10	10.85	10.87
標準偏差	1.24	1.24		1.25	1.24		1.21	1.24

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査