

表 2207 1セッションあたりの置換液量(L) Body Mass Index別（オフラインHDF 透析回数週3回 後希釈）

| Body Mass Index | 1未満 | 1～     | 5～     | 10～    | 15～   | 20～   | 合計      | 記載なし  | 総計    | 平均   | 標準偏差 |
|-----------------|-----|--------|--------|--------|-------|-------|---------|-------|-------|------|------|
| 14未満            |     | 5      | 15     |        |       | 1     | 21      |       | 21    | 7.00 | 3.44 |
| (%)             |     | (23.8) | (71.4) |        |       | (4.8) | (100.0) |       |       |      |      |
| 14～             |     | 23     | 90     | 42     | 1     |       | 156     | 5     | 161   | 7.22 | 2.39 |
| (%)             |     | (14.7) | (57.7) | (26.9) | (0.6) |       | (100.0) |       |       |      |      |
| 16～             |     | 44     | 282    | 169    | 2     |       | 497     | 28    | 525   | 7.90 | 2.19 |
| (%)             |     | (8.9)  | (56.7) | (34.0) | (0.4) |       | (100.0) |       |       |      |      |
| 18～             |     | 63     | 495    | 312    | 4     | 2     | 876     | 34    | 910   | 8.00 | 2.27 |
| (%)             |     | (7.2)  | (56.5) | (35.6) | (0.5) | (0.2) | (100.0) |       |       |      |      |
| 20～             |     | 70     | 444    | 329    | 1     | 1     | 845     | 28    | 873   | 8.06 | 2.27 |
| (%)             |     | (8.3)  | (52.5) | (38.9) | (0.1) | (0.1) | (100.0) |       |       |      |      |
| 22～             |     | 38     | 322    | 233    | 1     | 1     | 595     | 38    | 633   | 8.24 | 2.24 |
| (%)             |     | (6.4)  | (54.1) | (39.2) | (0.2) | (0.2) | (100.0) |       |       |      |      |
| 24～             |     | 23     | 211    | 144    | 2     | 1     | 381     | 13    | 394   | 8.27 | 2.25 |
| (%)             |     | (6.0)  | (55.4) | (37.8) | (0.5) | (0.3) | (100.0) |       |       |      |      |
| 26～             |     | 11     | 93     | 82     |       |       | 186     | 11    | 197   | 8.33 | 2.10 |
| (%)             |     | (5.9)  | (50.0) | (44.1) |       |       | (100.0) |       |       |      |      |
| 28～             |     | 10     | 49     | 38     |       |       | 97      | 9     | 106   | 8.02 | 2.23 |
| (%)             |     | (10.3) | (50.5) | (39.2) |       |       | (100.0) |       |       |      |      |
| 30～             |     | 6      | 50     | 34     |       |       | 90      | 1     | 91    | 8.30 | 2.11 |
| (%)             |     | (6.7)  | (55.6) | (37.8) |       |       | (100.0) |       |       |      |      |
| 合計              |     | 293    | 2,051  | 1,383  | 11    | 6     | 3,744   | 167   | 3,911 | 8.05 | 2.26 |
| (%)             |     | (7.8)  | (54.8) | (36.9) | (0.3) | (0.2) | (100.0) |       |       |      |      |
| 記載なし            |     | 33     | 143    | 88     |       | 1     | 265     | 18    | 283   | 7.58 | 2.37 |
| (%)             |     | (12.5) | (54.0) | (33.2) |       | (0.4) | (100.0) |       |       |      |      |
| 総計              |     | 326    | 2,194  | 1,471  | 11    | 7     | 4,009   | 185   | 4,194 | 8.02 | 2.27 |
| (%)             |     | (8.1)  | (54.7) | (36.7) | (0.3) | (0.2) | (100.0) |       |       |      |      |
| 平均              |     | 20.67  | 21.07  | 21.46  | 19.80 | 20.19 | 21.18   | 21.43 | 21.19 |      |      |
| 標準偏差            |     | 3.78   | 3.64   | 3.50   | 3.29  | 4.04  | 3.60    | 3.57  | 3.60  |      |      |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査