

表 2208 1セッションあたりの置換液量(L) 透析前BUN(mg/dL)別（オフラインHDF 透析回数週3回 透析歴2年以上 後希釈）

透析前BUN	1未満	1～	5～	10～	15～	20～	合計	記載なし	総計	平均	標準偏差
20未満		1	5	1	1		8		8	7.50	4.75
(%)		(12.5)	(62.5)	(12.5)	(12.5)		(100.0)				
20～		5	22	13			40	1	41	7.67	2.08
(%)		(12.5)	(55.0)	(32.5)			(100.0)				
30～		10	117	50			177	11	188	7.77	1.93
(%)		(5.6)	(66.1)	(28.2)			(100.0)				
40～		48	266	198	2	3	517	26	543	8.03	2.51
(%)		(9.3)	(51.5)	(38.3)	(0.4)	(0.6)	(100.0)				
50～		73	487	354	2		916	30	946	8.11	2.19
(%)		(8.0)	(53.2)	(38.6)	(0.2)		(100.0)				
60～		61	497	368	2	2	930	43	973	8.19	2.27
(%)		(6.6)	(53.4)	(39.6)	(0.2)	(0.2)	(100.0)				
70～		46	326	253		2	627	37	664	8.15	2.24
(%)		(7.3)	(52.0)	(40.4)		(0.3)	(100.0)				
80～		22	163	97	1		283	9	292	7.93	2.17
(%)		(7.8)	(57.6)	(34.3)	(0.4)		(100.0)				
90～		11	61	43			115	4	119	8.00	2.22
(%)		(9.6)	(53.0)	(37.4)			(100.0)				
100～		4	22	7			33	3	36	7.52	2.05
(%)		(12.1)	(66.7)	(21.2)			(100.0)				
110～			3				3		3	6.67	1.15
(%)			(100.0)				(100.0)				
120～		1	2	1			4		4	6.50	2.52
(%)		(25.0)	(50.0)	(25.0)			(100.0)				
合計		282	1,971	1,385	8	7	3,653	164	3,817	8.08	2.26
(%)		(7.7)	(54.0)	(37.9)	(0.2)	(0.2)	(100.0)				
記載なし		9	32	15			56	6	62	7.29	2.33
(%)		(16.1)	(57.1)	(26.8)			(100.0)				
総計		291	2,003	1,400	8	7	3,709	170	3,879	8.07	2.26
(%)		(7.8)	(54.0)	(37.7)	(0.2)	(0.2)	(100.0)				
平均		61.47	61.78	62.03	52.88	57.57	61.83	61.88	61.83		
標準偏差		16.62	15.67	14.59	22.87	13.29	15.36	15.29	15.35		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査