

表 2233 1セッションあたりの置換液量(L) 原疾患別 (オンラインHDF 透析回数週3回 後希釈)

原疾患	1未満	1～	5～	10～	15～	20～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎		33	159	240	60	37	529	11	540	10.70	4.70
(%)		(6.2)	(30.1)	(45.4)	(11.3)	(7.0)	(100.0)				
慢性腎盂腎炎		1	4	6	1	4	16		16	12.13	6.40
(%)		(6.3)	(25.0)	(37.5)	(6.3)	(25.0)	(100.0)				
急速進行性糸球体腎炎		1		3			4		4	8.50	4.43
(%)		(25.0)		(75.0)			(100.0)				
妊娠腎/妊娠中毒症		2	4	5	1		12		12	8.50	4.21
(%)		(16.7)	(33.3)	(41.7)	(8.3)		(100.0)				
その他の分類不能の腎炎			4	2		1	7		7	10.14	4.49
(%)			(57.1)	(28.6)		(14.3)	(100.0)				
多発性嚢胞腎		2	13	21	3	7	46	1	47	11.50	5.03
(%)		(4.3)	(28.3)	(45.7)	(6.5)	(15.2)	(100.0)				
腎硬化症		12	54	29	8	15	118	1	119	10.31	5.75
(%)		(10.2)	(45.8)	(24.6)	(6.8)	(12.7)	(100.0)				
悪性高血圧			3	3		2	8	1	9	12.38	5.40
(%)			(37.5)	(37.5)		(25.0)	(100.0)				
糖尿病性腎症		32	173	130	15	50	400	8	408	10.14	5.12
(%)		(8.0)	(43.3)	(32.5)	(3.8)	(12.5)	(100.0)				
SLE腎炎		1	4	4	3		12	2	14	10.75	4.65
(%)		(8.3)	(33.3)	(33.3)	(25.0)		(100.0)				
アミロイド腎			4	1		1	6		6	10.67	4.72
(%)			(66.7)	(16.7)		(16.7)	(100.0)				
痛風腎			2	3	3		8		8	12.00	4.11
(%)			(25.0)	(37.5)	(37.5)		(100.0)				
先天性代謝異常による腎不全			3				3		3	8.33	0.58
(%)			(100.0)				(100.0)				
腎・尿路結核			1				1		1	7.00	
(%)			(100.0)				(100.0)				
腎・尿路結石					1		1		1	16.00	
(%)					(100.0)		(100.0)				
腎・尿路腫瘍			2	1			3		3	9.00	1.00
(%)			(66.7)	(33.3)			(100.0)				
閉塞性尿路障害				2			2		2	11.00	1.41
(%)				(100.0)			(100.0)				
骨髄腫											
(%)											
腎形成不全				2			2	1	3	10.00	
(%)				(100.0)			(100.0)				
不明		1	36	22	6	17	82	1	83	11.73	5.65
(%)		(1.2)	(43.9)	(26.8)	(7.3)	(20.7)	(100.0)				
再導入		2	2	11	2	1	18		18	10.94	4.37
(%)		(11.1)	(11.1)	(61.1)	(11.1)	(5.6)	(100.0)				
その他			8	10	1	1	20	2	22	10.45	4.02
(%)			(40.0)	(50.0)	(5.0)	(5.0)	(100.0)				
合計		87	476	495	104	136	1,298	28	1,326	10.58	5.01
(%)		(6.7)	(36.7)	(38.1)	(8.0)	(10.5)	(100.0)				
記載なし											
(%)											
総計		87	476	495	104	136	1,298	28	1,326	10.58	5.01
(%)		(6.7)	(36.7)	(38.1)	(8.0)	(10.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査