

表 2301 身長(cm) 治療方法(新分類)別 (透析患者全体)

治療方法 (新分類)	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
施設血液透析	251	611	2,462	8,814	19,892	31,630	36,722	44,485	39,485	24,958	9,008	2,366	372	62	221,118	34,523	255,641	159.91	9.60
(%)	(0.1)	(0.3)	(1.1)	(4.0)	(9.0)	(14.3)	(16.6)	(20.1)	(17.9)	(11.3)	(4.1)	(1.1)	(0.2)	(0.0)	(100.0)				
Off-line HDF	10	12	53	227	532	884	976	1,023	960	682	258	69	15	2	5,703	612	6,315	159.88	9.77
(%)	(0.2)	(0.2)	(0.9)	(4.0)	(9.3)	(15.5)	(17.1)	(17.9)	(16.8)	(12.0)	(4.5)	(1.2)	(0.3)	(0.0)	(100.0)				
On-line HDF	30	58	243	937	2,508	4,412	5,245	6,273	5,885	4,194	1,562	436	65	8	31,856	4,234	36,090	160.91	9.49
(%)	(0.1)	(0.2)	(0.8)	(2.9)	(7.9)	(13.8)	(16.5)	(19.7)	(18.5)	(13.2)	(4.9)	(1.4)	(0.2)	(0.0)	(100.0)				
Push/Pull HDF			4	16	48	83	69	96	89	65	17	3			490	47	537	159.93	9.18
(%)			(0.8)	(3.3)	(9.8)	(16.9)	(14.1)	(19.6)	(18.2)	(13.3)	(3.5)	(0.6)			(100.0)				
ハイオフィルトレーション		2	4	23	28	55	39	60	48	29	11	4			303	38	341	158.41	10.15
(%)		(0.7)	(1.3)	(7.6)	(9.2)	(18.2)	(12.9)	(19.8)	(15.8)	(9.6)	(3.6)	(1.3)			(100.0)				
血液濾過	1	1		2	8	9	21	19	15	10	1				87	5	92	158.98	9.85
(%)	(1.1)	(1.1)		(2.3)	(9.2)	(10.3)	(24.1)	(21.8)	(17.2)	(11.5)	(1.1)				(100.0)				
血液吸着透析 (リクセル)		1	9	49	177	271	312	294	222	135	45	2			1,517	106	1,623	158.57	8.63
(%)		(0.1)	(0.6)	(3.2)	(11.7)	(17.9)	(20.6)	(19.4)	(14.6)	(8.9)	(3.0)	(0.1)			(100.0)				
血液吸着透析 (その他)						1	1		1	1	1				5	2	7	164.20	9.63
(%)						(20.0)	(20.0)		(20.0)	(20.0)	(20.0)				(100.0)				
在宅血液透析				2	10	23	38	70	71	96	50	12		1	373	148	521	166.47	8.32
(%)				(0.5)	(2.7)	(6.2)	(10.2)	(18.8)	(19.0)	(25.7)	(13.4)	(3.2)		(0.3)	(100.0)				
腹膜透析 (CAPD)	7	3	31	130	301	536	622	733	651	440	179	37	7	2	3,679	1,342	5,021	160.38	9.72
(%)	(0.2)	(0.1)	(0.8)	(3.5)	(8.2)	(14.6)	(16.9)	(19.9)	(17.7)	(12.0)	(4.9)	(1.0)	(0.2)	(0.1)	(100.0)				
腹膜透析 (APD)	48	4	29	77	123	235	329	432	408	306	127	32	4	2	2,156	683	2,839	160.11	14.33
(%)	(2.2)	(0.2)	(1.3)	(3.6)	(5.7)	(10.9)	(15.3)	(20.0)	(18.9)	(14.2)	(5.9)	(1.5)	(0.2)	(0.1)	(100.0)				
腹膜透析 (CCPD)	7	4	2	20	46	96	98	165	153	158	60	18	2		829	252	1,081	162.50	11.07
(%)	(0.8)	(0.5)	(0.2)	(2.4)	(5.5)	(11.6)	(11.8)	(19.9)	(18.5)	(19.1)	(7.2)	(2.2)	(0.2)		(100.0)				
合計	354	696	2,837	10,297	23,673	38,235	44,472	53,650	47,988	31,074	11,319	2,979	465	77	268,116	41,992	310,108	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.3)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし (%)																			
総計	354	696	2,837	10,297	23,673	38,235	44,472	53,650	47,988	31,074	11,319	2,979	465	77	268,116	41,992	310,108	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.3)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析 (CCPD) は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査