

表 2306 身長(cm) 原疾患別（透析患者全体）

原疾患	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	63	224	864	3,224	7,818	13,093	14,578	16,496	14,580	9,497	3,287	821	117	15	84,677	12,293	96,970	159.73	9.46
(%)	(0.1)	(0.3)	(1.0)	(3.8)	(9.2)	(15.5)	(17.2)	(19.5)	(17.2)	(11.2)	(3.9)	(1.0)	(0.1)	(0.0)	(100.0)				
慢性腎盂腎炎	3	14	33	155	337	503	479	445	345	208	85	26	4		2,637	405	3,042	157.59	9.68
(%)	(0.1)	(0.5)	(1.3)	(5.9)	(12.8)	(19.1)	(18.2)	(16.9)	(13.1)	(7.9)	(3.2)	(1.0)	(0.2)		(100.0)				
急速進行性糸球体腎炎	2	6	37	125	237	373	400	395	292	203	54	22	4		2,150	344	2,494	158.04	9.74
(%)	(0.1)	(0.3)	(1.7)	(5.8)	(11.0)	(17.3)	(18.6)	(18.4)	(13.6)	(9.4)	(2.5)	(1.0)	(0.2)		(100.0)				
妊娠腎/妊娠中毒症	1	2	16	93	300	489	379	151	41	10	1	1			1,484	130	1,614	152.92	6.19
(%)	(0.1)	(0.1)	(1.1)	(6.3)	(20.2)	(33.0)	(25.5)	(10.2)	(2.8)	(0.7)	(0.1)	(0.1)			(100.0)				
その他の分類不能の腎炎	7	1	18	43	108	188	199	212	212	142	58	17	2		1,207	166	1,373	159.71	10.94
(%)	(0.6)	(0.1)	(1.5)	(3.6)	(8.9)	(15.6)	(16.5)	(17.6)	(17.6)	(11.8)	(4.8)	(1.4)	(0.2)		(100.0)				
多発性嚢胞腎	10	7	46	219	712	1,409	1,732	1,694	1,605	1,380	667	190	45	4	9,720	1,286	11,006	161.59	9.77
(%)	(0.1)	(0.1)	(0.5)	(2.3)	(7.3)	(14.5)	(17.8)	(17.4)	(16.5)	(14.2)	(6.9)	(2.0)	(0.5)	(0.0)	(100.0)				
腎硬化症	30	122	465	1,416	2,671	3,673	4,322	5,090	3,895	2,143	653	150	17	1	24,648	3,650	28,298	158.22	9.67
(%)	(0.1)	(0.5)	(1.9)	(5.7)	(10.8)	(14.9)	(17.5)	(20.7)	(15.8)	(8.7)	(2.6)	(0.6)	(0.1)	(0.0)	(100.0)				
悪性高血圧	2	8	32	60	163	286	342	431	422	316	134	32	2	1	2,231	342	2,573	161.27	9.95
(%)	(0.1)	(0.4)	(1.4)	(2.7)	(7.3)	(12.8)	(15.3)	(19.3)	(18.9)	(14.2)	(6.0)	(1.4)	(0.1)	(0.0)	(100.0)				
糖尿病性腎症	126	167	762	3,136	7,477	12,554	15,680	21,427	20,282	13,439	5,018	1,377	214	51	101,710	16,371	118,081	161.21	9.46
(%)	(0.1)	(0.2)	(0.7)	(3.1)	(7.4)	(12.3)	(15.4)	(21.1)	(19.9)	(13.2)	(4.9)	(1.4)	(0.2)	(0.1)	(100.0)				
SLE腎炎	2	4	24	95	246	423	433	317	209	116	48	14	1		1,932	304	2,236	156.91	8.82
(%)	(0.1)	(0.2)	(1.2)	(4.9)	(12.7)	(21.9)	(22.4)	(16.4)	(10.8)	(6.0)	(2.5)	(0.7)	(0.1)		(100.0)				
アミロイド腎	1		10	34	52	76	64	56	41	30	9	1	1		375	76	451	156.03	9.94
(%)	(0.3)		(2.7)	(9.1)	(13.9)	(20.3)	(17.1)	(14.9)	(10.9)	(8.0)	(2.4)	(0.3)	(0.3)		(100.0)				
痛風腎			4	13	40	77	137	256	241	154	38	10	5		975	138	1,113	163.16	7.95
(%)			(0.4)	(1.3)	(4.1)	(7.9)	(14.1)	(26.3)	(24.7)	(15.8)	(3.9)	(1.0)	(0.5)		(100.0)				
先天性代謝異常による腎不全	3	4	10	5	30	29	48	41	42	23	7		1		243	29	272	157.61	11.73
(%)	(1.2)	(1.6)	(4.1)	(2.1)	(12.3)	(11.9)	(19.8)	(16.9)	(17.3)	(9.5)	(2.9)		(0.4)		(100.0)				
腎・尿路結核	1	3	5	7	27	29	35	41	19	10	2				179	27	206	156.22	9.15
(%)	(0.6)	(1.7)	(2.8)	(3.9)	(15.1)	(16.2)	(19.6)	(22.9)	(10.6)	(5.6)	(1.1)				(100.0)				
腎・尿路結石	4	2	6	25	53	87	95	106	71	39	17	4			509	72	581	158.13	9.59
(%)	(0.8)	(0.4)	(1.2)	(4.9)	(10.4)	(17.1)	(18.7)	(20.8)	(13.9)	(7.7)	(3.3)	(0.8)			(100.0)				
腎・尿路腫瘍		1	12	21	46	75	130	183	179	88	23	7			765	121	886	161.05	8.68
(%)		(0.1)	(1.6)	(2.7)	(6.0)	(9.8)	(17.0)	(23.9)	(23.4)	(11.5)	(3.0)	(0.9)			(100.0)				
閉塞性尿路障害	6	5	8	27	76	96	106	114	108	60	25	2	1		634	89	723	158.41	10.68
(%)	(0.9)	(0.8)	(1.3)	(4.3)	(12.0)	(15.1)	(16.7)	(18.0)	(17.0)	(9.5)	(3.9)	(0.3)	(0.2)		(100.0)				
骨髄腫	1	1	3	11	33	31	34	44	38	24	7	1	1		229	47	276	158.56	10.02
(%)	(0.4)	(0.4)	(1.3)	(4.8)	(14.4)	(13.5)	(14.8)	(19.2)	(16.6)	(10.5)	(3.1)	(0.4)	(0.4)		(100.0)				
腎形成不全	25	8	21	41	65	63	97	100	73	69	17	9			588	61	649	155.47	16.74
(%)	(4.3)	(1.4)	(3.6)	(7.0)	(11.1)	(10.7)	(16.5)	(17.0)	(12.4)	(11.7)	(2.9)	(1.5)			(100.0)				
不明	39	95	362	1,202	2,402	3,571	3,859	4,479	3,816	2,185	791	192	28	3	23,024	4,520	27,544	158.71	9.81
(%)	(0.2)	(0.4)	(1.6)	(5.2)	(10.4)	(15.5)	(16.8)	(19.5)	(16.6)	(9.5)	(3.4)	(0.8)	(0.1)	(0.0)	(100.0)				
再導入	3	6	18	34	130	216	305	382	386	254	109	33	3	1	1,880	283	2,163	161.78	9.50
(%)	(0.2)	(0.3)	(1.0)	(1.8)	(6.9)	(11.5)	(16.2)	(20.3)	(20.5)	(13.5)	(5.8)	(1.8)	(0.2)	(0.1)	(100.0)				
その他	25	16	81	311	650	894	1,018	1,190	1,091	684	269	70	19	1	6,319	1,238	7,557	159.34	10.49
(%)	(0.4)	(0.3)	(1.3)	(4.9)	(10.3)	(14.1)	(16.1)	(18.8)	(17.3)	(10.8)	(4.3)	(1.1)	(0.3)	(0.0)	(100.0)				
合計	354	696	2,837	10,297	23,673	38,235	44,472	53,650	47,988	31,074	11,319	2,979	465	77	268,116	41,992	310,108	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.3)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				
記載なし																			
(%)																			
総計	354	696	2,837	10,297	23,673	38,235	44,472	53,650	47,988	31,074	11,319	2,979	465	77	268,116	41,992	310,108	160.05	9.65
(%)	(0.1)	(0.3)	(1.1)	(3.8)	(8.8)	(14.3)	(16.6)	(20.0)	(17.9)	(11.6)	(4.2)	(1.1)	(0.2)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査